

2019 PRIOR AUTHORIZATION CRITERIA

TABLE OF CONTENTS

ABILIFY tablet.....	34
abiraterone 250 mg tablet.....	251
ABRAXANE	143
ABSTRAL.....	104
ACTIMMUNE	20
ADASUVE.....	34
ADCIRCA.....	229
ADEMPAS	231
AFINITOR	251
AFINITOR DISPERZ.....	251
AIMOVIG.....	21
ALDARA.....	141
ALECENSA.....	251
ALIMTA	143
ALIQOPA	143
ALUNBRIG.....	251
ambrisentan tablet	235
AMITIZA.....	24
amitriptyline tablet	131
AMOXAPINE.....	131
AMPYRA.....	87
ANADROL-50	27
ANDRODERM	32

ANDROGEL 1%.....	32
ANDROGEL 1.62%.....	32
ANDROXY	31
APOKYN	37
ARANESP.....	97
ARCALYST	38
aripiprazole ODT	34
aripiprazole solution, tablet	34
ARISTADA	34
ARISTADA INITIO	34
armodafinil tablet.....	39
ARZERRA.....	143
AUBAGIO.....	166
AURYXIA	42
AVASTIN.....	143
AVONEX	167
AVONEX PREFILLED KIT	167
AXIRON	32
BALVERSA	251
BAVENCIO	188
BELEODAQ	143
BENLYSTA	43
benztropine tablet	129
BESPONSA	143
BETASERON.....	168
bexarotene capsule.....	251

BLINCYTO	143
bosentan tablet	233
BOSULIF.....	251
BRAFTOVI	251
BUPHENYL tablet.....	288
buprenorphine transdermal patch	196
BUTRANS.....	196
CABOMETYX	251
CALQUENCE.....	251
CAPRELSA.....	251
CARBAGLU	76
CEREZYME	112
CHENODAL	77
CHLORPROMAZINE injection.....	34
chlorpromazine tablet.....	34
CHORIONIC GONADOTROPIN.....	78
cinacalcet tablet	79
CINRYZE	121
CLEMASTINE tablet	129
clobazam suspension, tablet.....	53
clomipramine capsule	131
clonazepam ODT	45
clonazepam tablet.....	45
clorazepate tablet.....	47
clozapine ODT 12.5 mg, 25 mg, 100 mg	34
clozapine tablet.....	34

CLOZARIL	34
COMETRIQ.....	251
COPAXONE.....	170
COPIKTRA.....	251
CORLANOR.....	80
COSENTYX	55
COTELLIC	251
CRESEMBA.....	81
CRYSVITA	82
cyclobenzaprine tablet	130
CYRAMZA	143
CYSTAGON.....	84
DAKLINZA	85
dalfampridine ER tablet.....	87
DALIRESP	88
danazol capsule	25
DARZALEX.....	143
DAURISMO.....	251
deferasirox tablet	148
DEPO-TESTOSTERONE	28
desipramine tablet.....	131
DIAZEPAM 1 mg/mL oral solution	49
diazepam intensol concentrate	49
diazepam tablet.....	49
diclofenac 1% gel.....	282
dicyclomine capsule, tablet.....	129

doxepin capsule, oral concentrate	131
doxorubicin liposomal injection	143
DUPIXENT	89
EGRIFTA	91
ELELYSO.....	114
ELIDEL.....	40
EMGALITY	92
EMPLICITI	143
ENBREL.....	57
EPCLUSA	94
EPIDIOLEX.....	96
EPOGEN.....	99
ERBITUX	143
ERIVEDGE	251
ERLEADA	251
erlotinib tablet.....	251
ESBRIET.....	145
EXJADE	148
FANAPT	34
FANAPT TITRATION PACK.....	34
FARYDAK.....	251
FAZACLO ODT 25 mg, 100 mg.....	34
fentanyl citrate oral lozenge	105
fentanyl transdermal patch.....	198
FIRAZYR.....	123
FLUPHENAZINE concentrate, elixir, injection	34

fluphenazine decanoate 25 mg/mL injection.....	34
fluphenazine tablet.....	34
FOLOTYN.....	143
FORTEO.....	106
GAMMAGARD.....	108
GAMMAGARD SD.....	108
GAMMAPLEX.....	73
GAMUNEX-C.....	108
GATTEX.....	111
GAZYVA.....	143
GEODON.....	34
GILENYA 0.5 mg capsule.....	169
GILOTRIF.....	251
glatiramer injection.....	170
GLEEVEC.....	251
GRASTEK.....	210
H.P. ACTHAR GEL.....	18
HAEGARDA.....	125
HALAVEN.....	143
HALDOL.....	34
HALDOL DECANOATE.....	34
haloperidol concentrate, injection, tablet.....	34
haloperidol decanoate injection.....	34
HARVONI.....	127
HERCEPTIN.....	143
HERCEPTIN HYLECTA.....	143

HETLIOZ.....	128
HEXALEN	251
HUMIRA.....	59
HUMIRA KIT	59
hydroxyzine hcl syrup, tablet.....	129
IBRANCE	251
icatibant 30 mg/3 mL injection	123
ICLUSIG.....	251
IDHIFA	251
ILARIS.....	139
imatinib mesylate tablet	251
IMBRUVICA	251
IMFINZI	189
imipramine hcl tablet.....	131
imiquimod 5% cream	141
INBRIJA	142
INLYTA	251
INVEGA	34
INVEGA SUSTENNA.....	34
INVEGA TRINZA	34
IRESSA.....	251
ISTODAX	143
JADENU.....	149
JADENU SPRINKLE.....	149
JAKAFI.....	251
JEVTANA.....	143

JUXTAPID.....	135
KADCYLA	143
KALYDECO	150
KANJINTI	143
KEYTRUDA	190
KINERET	61
KISQALI	251
KISQALI FEMARA CO PACK.....	251
KORLYM.....	152
KUVAN.....	153
KYNAMRO.....	137
KYPROLIS	143
LARTRUVO	143
LATUDA.....	34
LAZANDA	103
LENVIMA	251
LETAIRIS	235
LIBTAYO.....	191
lidocaine 2% gel.....	154
lidocaine 4% solution	157
lidocaine 5% ointment.....	155
lidocaine 5% transdermal patch.....	156
lidocaine/prilocaine 2.5-2.5% cream	158
LIDODERM.....	156
linezolid suspension, tablet.....	159
LINZESS.....	161

LONSURF	251
lorazepam tablet	51
LORBRENA	251
loxapine capsule	34
LUMOXITI	143
LYNPARZA	251
MATULANE	251
MAVYRET	162
MAYZENT	171
megestrol suspension, tablet	131
MEKINIST	251
MEKTOVI	251
memantine solution, tablet	164
memantine titration pak.....	164
METHYLTESTOSTERONE 10 mg capsule.....	31
miglustat 100 mg capsule	274
modafinil tablet.....	165
MOLINDONE	34
morphine sulfate ER tablet.....	202
MVASI	143
MYALEPT	176
MYLOTARG	143
NAMENDA	164
NAMENDA TITRATION PAK.....	164
NATPARA	178
NERLYNX	251

NEXAVAR.....	251
NINLARO	251
NORPRAMIN	131
NORTHERA.....	180
nortriptyline capsule	131
NORTRIPTYLINE solution.....	131
NOXAFIL.....	182
NUCYNTA ER.....	206
NUEDEXTA	184
NUPLAZID	185
NUVIGIL.....	39
OICALIVA	186
octreotide injection	260
ODOMZO.....	251
OFEV	146
olanzapine injection, tablet.....	34
olanzapine ODT	34
OMNITROPE	118
ONFI	53
ONIVYDE	143
OPDIVO	192
OPSUMIT.....	237
ORALAIR	212
ORENCIA.....	63
ORKAMBI	216
OTEZLA	218

oxandrolone tablet	26
OXYCONTIN.....	204
paliperidone ER tablet.....	34
PALYNZIQ	220
paroxetine tablet	131
PAXIL.....	131
PEGASYS.....	222
PEGASYS PROCLICK	222
PERJETA.....	143
perphenazine tablet	34
PERSERIS.....	34
pimecrolimus 1% cream.....	40
PIQRAY	252
PLEGRIDY	172
PLEGRIDY STARTER PACK	172
POLIVY	143
POMALYST	252
PORTRAZZA	143
posaconazole 100 mg delayed release tablet.....	182
POTELIGEO	143
PREGNYL.....	78
PROCRIT.....	99
PROLASTIN-C.....	23
PROLIA.....	223
PROMACTA.....	226
promethazine 12.5 mg, 25 mg suppository	129

promethazine syrup, tablet.....	129
protriptyline tablet.....	131
PRUDOXIN	281
quetiapine ER tablet.....	34
quetiapine tablet.....	34
RAGWITEK.....	214
REGRANEX.....	245
RELISTOR	246
RENFLEXIS	65
REPATHA	247
RESTASIS	195
RESTASIS MULTIDOSE	195
RETACRIT	101
REVLIMID	252
REXULTI.....	34
RISPERDAL.....	34
RISPERDAL CONSTA.....	34
RISPERDAL M-TAB 2 mg	34
risperidone ODT.....	34
risperidone solution, tablet.....	34
RITUXAN	67
RITUXAN HYCELA.....	69
RUBRACA	252
RYDAPT	252
SAMSCA.....	250
SANDOSTATIN LAR	262

SAPHRIS	34
SENSIPAR	79
SEROQUEL	34
SEROQUEL XR	34
SIGNIFOR.....	256
SIGNIFOR LAR.....	254
sildenafil 20 mg tablet	239
SIVEXTRO tablet	258
sodium phenylbutyrate powder, tablet	288
SOMATULINE DEPOT	264
SOMAVERT	266
SOVALDI	268
SPRAVATO	270
SPRYCEL	252
STELARA.....	71
STIVARGA.....	252
STRENSIQ.....	272
SURMONTIL.....	131
SUTENT	252
SYLATRON.....	252
SYMDEKO	276
SYMPAZAN	54
SYNRIBO.....	143
SYPRINE	284
tacrolimus ointment.....	41
tadalafil 20 mg tablet (generic for Adcirca)	229

TAFINLAR.....	252
TAGRISSE	252
TALZENNA	252
TARCEVA.....	252
TARGRETIN 75 mg capsule.....	252
TASIGNA	252
TECENTRIQ	193
TECFIDERA.....	173
TECFIDERA STARTER PACK.....	173
TECHNIVIE.....	278
testosterone 1% gel	32
testosterone 1.62% gel	32
testosterone 30 mg/actuation solution	32
testosterone cypionate injection.....	28
testosterone enanthate injection	30
tetrabenazine tablet	280
THALOMID	252
thioridazine tablet.....	35
thiothixene capsule	35
TIBSOVO	252
TRACLEER.....	233
tramadol ER tablet	208
TRELSTAR	283
TRELSTAR MIXJECT.....	283
tretinoin capsule.....	252
trientine capsule.....	284

trifluoperazine tablet.....	35
trimipramine capsule.....	131
TYKERB.....	252
TYMLOS	286
TYSABRI.....	174
UNITUXIN	143
UPTRAVI	241
VECTIBIX.....	143
VELCADE	143
VENCLEXTA.....	252
VENCLEXTA STARTING PACK.....	252
VENTAVIS	243
VERSACLOZ	35
VERZENIO.....	252
VFEND IV	293
VIEKIRA PAK.....	289
VIEKIRA XR.....	291
VITRAKVI.....	252
VIZIMPRO.....	252
VOLTAREN 1% gel.....	282
voriconazole injection, suspension, tablet.....	293
VOSEVI.....	295
VOTRIENT	252
VPRIV	116
VRAYLAR	35
VYNDAMAX.....	296

VYNDAQEL	297
VYXEOS	143
XALKORI	252
XENAZINE	280
XGEVA.....	298
XOLAIR.....	300
XOSPATA	252
XPOVIO	252
XTANDI.....	252
XYREM	303
YERVOY	194
YONDELIS.....	143
YONSA	252
zaleplon capsule	133
ZALTRAP	144
ZEJULA.....	252
ZELBORAF	252
ZEPATIER	304
ziprasidone capsule	35
ZOHYDRO ER	200
ZOLINZA.....	252
zolpidem tablet.....	134
ZONALON.....	281
ZYDELIG.....	252
ZYKADIA.....	252
ZYPREXA	35

ZYPREXA RELPREVV	35
ZYPREXA ZYDIS.....	35
ZYTIGA	252
ZYVOX suspension, tablet.....	159

Prior Authorization Group – Acthar HP Gel PA

Drug Name(s):

H.P. ACTHAR GEL

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. ONE of the following:

- a. Patient has a diagnosis of infantile spasm OR
- b. Patient has a diagnosis of nephrotic syndrome AND ONE of the following:
 - i. Patient has failed a conventional agent (i.e. prednisone, cyclophosphamide, tacrolimus) for the requested indication OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a conventional agent OR
- c. Patient has a diagnosis of multiple sclerosis AND ALL of the following:
 - i. Patient is experiencing an acute exacerbation AND
 - ii. If indicated, there is evidence of a claim that the patient is currently on a disease modifying drug (DMD) within the past 90 days (e.g. Aubagio, Avonex, Betaseron, Extavia, Gilenya, glatiramer (i.e. Copaxone, Glatopa), Lemtrada, mitoxantrone, Ocrevus, Plegridy, Rebif, Tecfidera, or Tysabri) to control disease progression OR has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a DMD AND
 - iii. Patient has failed corticosteroid therapy within the last 30 days or has an FDA labeled contraindication to corticosteroid therapy (e.g. methylprednisolone 1gm IV for 3-5 days) OR

Criteria continues: see Other Criteria

Age Restrictions:

For diagnosis of infantile spasm, patient is less than 24 months of age. For diagnosis of nephrotic syndrome, patient is greater than 2 years of age.

Prescriber Restrictions:

Coverage Duration:

6 months for infantile spasm, 1 month for all other indications

Other Criteria:

- d. Patient has a diagnosis of rheumatic disorder (e.g. ankylosing spondylitis, juvenile idiopathic arthritis, juvenile rheumatoid arthritis, psoriatic arthritis, rheumatoid arthritis) AND BOTH of the following:
 - i. If indicated, there is evidence of a claim that the patient is currently being treated with a conventional agent within the past 90 days (e.g. DMARD [methotrexate, leflunomide], biologics [Humira, Enbrel]) to control disease progression OR has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a conventional agent AND
 - ii. Patient has failed corticosteroid therapy within the last 30 days or has an FDA labeled contraindication to corticosteroid therapy (e.g. methylprednisolone 1gm IV for 3-5 days) OR
 - e. Patient has a diagnosis of systemic lupus erythematosus (SLE) disease AND ALL of the following:
 - i. Patient has a history of positive antinuclear antibody (ANA) and/or positive anti-dsDNA results AND
 - ii. Patient has a history of 3 other SLE diagnostic criteria (i.e. malar rash, discoid rash, photosensitivity, oral ulcers, nonerosive arthritis, serositis [e.g. pleuritis/pericarditis], renal disorder [e.g. persistent proteinuria greater than 0.5 grams/day or cellular casts], hematologic disorder [e.g. hemolytic anemia (with reticulocytosis), leukopenia, lymphopenia, or thrombocytopenia], and/or immunologic disorder [e.g. positive finding of antiphospholipid antibodies or anti-Sm antibodies]) AND
 - iii. ONE of the following:
 - 1. There is evidence of a claim that the patient is currently on a SLE treatment regimen within the past 90 days comprised of at least ONE of the following: corticosteroids (e.g. methylprednisolone, prednisone), antimalarials (e.g. hydroxychloroquine, chloroquine), prescription nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g. ibuprofen, naproxen), and/or immunosuppressive (e.g. azathioprine, methotrexate, cyclosporine, or oral cyclophosphamide) OR
 - 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to any standard of care drug class listed above OR
 - f. Patient has another FDA approved indication AND
 - i. Patient has failed corticosteroid therapy within the last 30 days or has an FDA labeled contraindication to corticosteroids therapy (e.g. methylprednisolone 1gm IV for 3-5 days) OR
 - g. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
 - i. Patient has failed corticosteroid therapy within the last 30 days or has an FDA labeled contraindication to corticosteroids therapy (e.g. methylprednisolone 1gm IV for 3-5 days) AND
2. The dose is within the FDA labeled or CMS approved compendia dosing for the requested indication

Prior Authorization Group – Actimmune PA

Drug Name(s):

ACTIMMUNE

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.

Criteria for approval require BOTH of the following:

1. ONE of the following:

- a. Patient has an FDA labeled indication for the requested agent OR
- b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. The dose requested is within the FDA labeled or CMS approved compendia dosing for the requested indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Aimovig PA

Drug Name(s):

AIMOVIG

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of migraine AND
2. Patient has 4 migraine headaches or more per month AND
3. ONE of the following:
 - a. Patient has failed at least TWO conventional migraine prophylaxis agents from TWO different classes (e.g. beta blockers [propranolol], anticonvulsants [divalproex, topiramate]) OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to TWO conventional migraine prophylaxis agents from TWO different classes AND
4. ONE of the following:
 - a. Patient is NOT currently taking another calcitonin gene-related peptide (CGRP) agent OR
 - b. Patient is currently being treated with another calcitonin gene-related peptide (CGRP) agent AND will discontinue prior to initiating therapy with the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's prior authorization criteria AND
2. Patient has a diagnosis of migraine AND
3. Patient has shown clinical benefit with the requested agent AND
4. ONE of the following:
 - a. Patient is NOT currently taking another calcitonin gene-related peptide (CGRP) agent OR
 - b. Patient is currently being treated with another calcitonin gene-related peptide (CGRP) agent AND will discontinue prior to continuing therapy with the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Alpha-1-Proteinase Inhibitor PA - Prolastin-C

Drug Name(s):

PROLASTIN-C

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of alpha-1 antitrypsin deficiency (AATD) AND
2. Patient has a pre-treatment serum alpha-1 antitrypsin (AAT) level less than 11 µM/L (80 mg/dL by immunodiffusion or 57 mg/dL using nephelometry) AND
3. Patient has one of the following phenotype variants associated with AATD: PiZZ, PiSZ, PiZ/Null or PiNull/Null AND
4. Patient has emphysema with a documented baseline FEV1 of 65% or less of predicted AND
5. The dose requested is within the FDA labeled dose

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of alpha-1 antitrypsin deficiency (AATD) with clinically evident emphysema AND
3. Patient has shown clinical benefit with the requested agent AND
4. The dose requested is within the FDA labeled dose

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Amitiza PA

Drug Name(s):

AMITIZA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. Patient has ONE of the following diagnoses:

A. Chronic idiopathic constipation with documentation of symptoms for at least 3 months OR

B. Irritable bowel syndrome with constipation with documentation of symptoms for at least 3 months AND the patient is female OR

C. Opioid-induced constipation with chronic non-cancer pain, including patients with chronic pain related to prior cancer or its treatment AND BOTH of the following:

i. Patient has chronic use of an opioid agent within the past 90 days AND

ii. Patient has NOT received methadone within the past 90 days AND

2. ONE of the following:

A. Patient has a medication history that includes lactulose or polyethylene glycol 3350 OR

B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to lactulose or polyethylene glycol 3350

Age Restrictions:

Patient is 18 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Anabolic Steroid PA - Danazol

Drug Name(s):

danazol capsule

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. Patient has ONE of the following diagnoses:
 - a. Patient has an FDA labeled indication for the requested agent OR
 - b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - a. Patient is NOT currently treated with another androgen or anabolic steroid within the past 90 days OR
 - b. The current androgen or anabolic steroid will be discontinued prior to starting the requested agent OR
 - c. Prescriber has submitted documentation in support of therapy with more than one agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Anabolic Steroid PA - Oxandrolone

Drug Name(s):

oxandrolone tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. Patient has ONE of the following diagnoses:
 - a. Patient has AIDS/HIV-associated wasting syndrome (defined as unexplained involuntary weight loss greater than 10% baseline body weight with obvious wasting or body mass index less than 18.5 kg/m²) AND all other causes of weight loss have been ruled out OR
 - b. Patient is a female child or adolescent with Turner syndrome AND is currently receiving growth hormone OR
 - c. Patient has weight loss following extensive surgery, chronic infections, or severe trauma OR
 - d. Patient has chronic pain from osteoporosis OR
 - e. Patient is on long-term administration of oral or injectable corticosteroids AND
2. ONE of the following:
 - a. Patient is NOT currently treated with another androgen or anabolic steroid within the past 90 days OR
 - b. The current androgen or anabolic steroid will be discontinued prior to starting the requested agent OR
 - c. Prescriber has submitted documentation in support of therapy with more than one agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Anabolic Steroid PA - Oxymetholone

Drug Name(s):

ANADROL-50

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:
 - a. Patient has anemia caused by deficient red cell production, including acquired aplastic anemia, congenital aplastic anemia, myelofibrosis and the hypoplastic anemias due to the administration of myelotoxic drugs OR
 - b. Patient has anemia associated with chronic renal failure AND ONE of the following:
 - i. Patient's medication history indicates previous use of an erythropoiesis-stimulating agent OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication or hypersensitivity to an erythropoiesis-stimulating agent AND
2. Patient has a hematocrit (Hct) value less than 30% AND
3. ONE of the following:
 - a. Patient is NOT currently treated with another androgen or anabolic steroid within the past 90 days OR
 - b. The current androgen or anabolic steroid will be discontinued prior to starting the requested agent OR
 - c. Prescriber has submitted documentation in support of therapy with more than one agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Androgen Injectable PA - testosterone cypionate

Drug Name(s):

DEPO-TESTOSTERONE

testosterone cypionate injection

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:
 - a. Patient is a male with AIDS/HIV-associated wasting syndrome, defined as unexplained involuntary weight loss (greater than 10% baseline body weight) with obvious wasting OR body mass index less than 18.5 kg/m² AND all other causes of weight loss have been ruled out OR
 - b. Patient is a female with metastatic/inoperable breast cancer OR
 - c. Patient is a male with primary or secondary (hypogonadotropic) hypogonadism OR
 - d. Patient is an adolescent male with delayed puberty AND
2. Males only: Patient has a measured current or pretreatment total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR a free serum testosterone level that is below the testing laboratory's lower limit of the normal range AND
3. ONE of the following:
 - a. Patient is NOT currently treated with another androgen or anabolic steroid within the past 90 days OR
 - b. The current androgen or anabolic steroid will be discontinued prior to starting the requested agent OR
 - c. Prescriber has submitted documentation in support of therapy with more than one agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for delayed puberty, 12 months for all other indications

Other Criteria:

Prior Authorization Group – Androgen Injectable PA - testosterone enanthate

Drug Name(s):

testosterone enanthate injection

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:
 - a. Patient is a male with AIDS/HIV-associated wasting syndrome, defined as unexplained involuntary weight loss (greater than 10% baseline body weight) with obvious wasting OR body mass index less than 18.5 kg/m² AND all other causes of weight loss have been ruled out OR
 - b. Patient is a female with metastatic/inoperable breast cancer OR
 - c. Patient is a male with primary or secondary (hypogonadotropic) hypogonadism OR
 - d. Patient is an adolescent male with delayed puberty AND
2. Males only: Patient has a measured current or pretreatment total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR a free serum testosterone level that is below the testing laboratory's lower limit of the normal range AND
3. ONE of the following:
 - a. Patient is NOT currently treated with another androgen or anabolic steroid within the past 90 days OR
 - b. The current androgen or anabolic steroid will be discontinued prior to starting the requested agent OR
 - c. Prescriber has submitted documentation in support of therapy with more than one agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for delayed puberty, 12 months for all other indications

Other Criteria:

Prior Authorization Group – Androgen Oral PA

Drug Name(s):

ANDROXY

METHYLTESTOSTERONE 10 mg capsule

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:
 - a. Patient is a male with cryptorchidism OR
 - b. Patient is a male with hypogonadism OR
 - c. Patient is an adolescent male with delayed puberty OR
 - d. Patient is a female with metastatic/inoperable breast cancer AND
2. Males only: Patient has a measured current or pretreatment total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR a free serum testosterone level that is below the testing laboratory's lower limit of the normal range AND
3. ONE of the following:
 - a. Patient is NOT currently treated with another androgen or anabolic steroid within the past 90 days OR
 - b. The current androgen or anabolic steroid will be discontinued prior to starting the requested agent OR
 - c. Prescriber has submitted documentation in support of therapy with more than one agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 months for delayed puberty, 12 months for all other indications

Other Criteria:

Prior Authorization Group – Androgen Topical PA

Drug Name(s):

ANDRODERM

ANDROGEL 1%

ANDROGEL 1.62%

AXIRON

testosterone 1% gel

testosterone 1.62% gel

testosterone 30 mg/actuation solution

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:
 - a. Patient has AIDS/HIV-associated wasting syndrome, defined as unexplained involuntary weight loss (greater than 10% baseline body weight) with obvious wasting OR body mass index less than 18.5 kg/m² AND all other causes of weight loss have been ruled out OR
 - b. Patient is a male with primary or secondary (hypogonadotropic) hypogonadism AND
2. Males only: Patient has a measured current or pretreatment total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR a free serum testosterone level that is below the testing laboratory's lower limit of the normal range AND
3. ONE of the following:
 - a. Patient is NOT currently treated with another androgen or anabolic steroid within the past 90 days OR
 - b. The current androgen or anabolic steroid will be discontinued prior to starting the requested agent OR
 - c. Prescriber has submitted documentation in support of therapy with more than one agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Antipsychotics PA

Drug Name(s):

ABILIFY tablet
ADASUVE
aripiprazole ODT
aripiprazole solution, tablet
ARISTADA
ARISTADA INITIO
CHLORPROMAZINE injection
chlorpromazine tablet
clozapine ODT 12.5 mg, 25 mg, 100 mg
clozapine tablet
CLOZARIL
FANAPT
FANAPT TITRATION PACK
FAZACLO ODT 25 mg, 100 mg
FLUPHENAZINE concentrate, elixir, injection
fluphenazine decanoate 25 mg/mL injection
fluphenazine tablet
GEODON
HALDOL
HALDOL DECANOATE
haloperidol concentrate, injection, tablet
haloperidol decanoate injection
INVEGA
INVEGA SUSTENNA
INVEGA TRINZA
LATUDA
loxapine capsule
MOLINDONE
olanzapine injection, tablet
olanzapine ODT
paliperidone ER tablet
perphenazine tablet
PERSERIS
quetiapine ER tablet
quetiapine tablet
REXULTI
RISPERDAL
RISPERDAL CONSTA
RISPERDAL M-TAB 2 mg
risperidone ODT
risperidone solution, tablet
SAPHRIS
SEROQUEL
SEROQUEL XR

thioridazine tablet
thiothixene capsule
trifluoperazine tablet
VERSACLOZ
VRAYLAR
ziprasidone capsule
ZYPREXA
ZYPREXA RELPREVV
ZYPREXA ZYDIS

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only. PA does NOT apply to patients less than 65 years of age.

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently using the requested agent OR
 - C. IF dementia-related psychosis, BOTH of the following:
 - i. Dementia-related psychosis is determined to be severe or the associated agitation, combativeness, or violent behavior puts the patient or others in danger AND
 - ii. Prescriber has documented that s/he has discussed the risk of increased mortality with the patient and/or the patient's surrogate decision maker

Approval authorizations will apply to the requested medication only.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Apokyn PA

Drug Name(s):

APOKYN

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Receiving a 5-HT3 antagonist (e.g., ondansetron, granisetron, dolasetron, palonosetron, alosetron) concomitantly with the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. The requested agent will be used to treat acute, intermittent hypomobility, “off” episodes (muscle stiffness, slow movements, or difficulty starting movement) associated with advanced Parkinson’s disease AND
2. There is evidence of a claim that the patient is receiving concurrent therapy for Parkinson’s disease (e.g., levodopa, dopamine agonist, or monoamine oxidase B inhibitor) within the past 30 days

Age Restrictions:

Prescriber Restrictions:

Prescriber is a neurologist or the prescriber has consulted with a neurologist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Arcalyst PA

Drug Name(s):

ARCALYST

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

An active or chronic infection (e.g. tuberculosis, HIV, hepatitis B/C)

Required Medical:

Criteria for approval require BOTH of the following:

1. Patient has been diagnosed with Cryopyrin-Associated Periodic Syndromes (CAPS) including Familial Cold Auto-inflammatory Syndrome (FCAS) or Muckle-Wells Syndrome (MWS) AND
2. ONE of the following:
 - A. Patient is NOT currently being treated with another biologic agent OR
 - B. Patient is currently being treated with another biologic agent and the agent will be discontinued prior to initiating the requested agent

Age Restrictions:

Patient is at least 12 years of age

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Armodafinil PA

Drug Name(s):

**armodafinil tablet
NUVIGIL**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require BOTH of the following:

1. ONE of the following:

- A. Patient has an FDA labeled indication for the requested agent OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. Patient is receiving only one of the listed agents, armodafinil OR modafinil, within the past 90 days OR
- B. Patient has been treated with modafinil within the past 90 days AND will discontinue prior to starting the requested agent

Age Restrictions:

Patient is 17 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Atopic Dermatitis PA - Elidel

Drug Name(s):

ELIDEL

pimecrolimus 1% cream

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require ONE of the following:

1. Patient has a diagnosis of atopic dermatitis or vulvar lichen sclerosus AND ONE of the following:
 - A. Patient has had a trial and failure of a topical corticosteroid or topical corticosteroid combination preparation (e.g. clobetasol, hydrocortisone, triamcinolone) OR
 - B. Patient has a documented intolerance, FDA labeled contraindication or hypersensitivity to a topical corticosteroid or topical corticosteroid combination preparation OR
2. Patient has a diagnosis of facial seborrheic dermatitis associated with HIV infection AND BOTH of the following:
 - A. Patient is currently on an antiretroviral treatment regimen AND
 - B. ONE of the following:
 - i. Patient has had a trial and failure of a topical corticosteroid or topical antifungal treatment (e.g. hydrocortisone, triamcinolone, ketoconazole, nystatin-triamcinolone) OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication or hypersensitivity to a topical corticosteroid or topical antifungal treatment OR
3. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Atopic Dermatitis PA - Tacrolimus

Drug Name(s):

tacrolimus ointment

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require ONE of the following:

1. Patient has a diagnosis of atopic dermatitis AND ONE of the following:
 - A. Patient has had a trial and failure with a topical corticosteroid or topical corticosteroid combination preparation (e.g. clobetasol, hydrocortisone, triamcinolone) OR
 - B. Patient has a documented intolerance, FDA labeled contraindication or hypersensitivity to a topical corticosteroid or topical corticosteroid combination preparation OR
2. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Auryxia PA

Drug Name(s):

AURYXIA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Requested agent will be used as iron replacement therapy to treat iron deficiency anemia AND FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require the following:

1. Patient has a diagnosis of hyperphosphatemia in chronic kidney disease on dialysis

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Benlysta PA

Drug Name(s):

BENLYSTA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of active systemic lupus erythematosus (SLE) disease AND
2. Patient has a history of positive antinuclear antibody (ANA) and/or positive anti-dsDNA results AND
3. Patient has a history of 3 other SLE diagnostic criteria (i.e. malar rash, discoid rash, photosensitivity, oral ulcers, nonerosive arthritis, serositis [e.g. pleuritis/pericarditis], renal disorder [e.g. persistent proteinuria greater than 0.5 grams/day or cellular casts], hematologic disorder [e.g. hemolytic anemia (with reticulocytosis), leukopenia, lymphopenia, or thrombocytopenia], and/or immunologic disorder [e.g. positive finding of antiphospholipid antibodies or anti-Sm antibodies]) AND
4. ONE of the following:
 - A. There is evidence of a claim that the patient is currently on SLE treatment regimen within the past 90 days comprised of at least ONE of the following: corticosteroids (e.g. methylprednisolone, prednisone), antimalarials (e.g. hydroxychloroquine, chloroquine), prescription nonsteroidal anti-inflammatory drugs (NSAIDS) (e.g. ibuprofen, naproxen), and/or immunosuppressives (e.g. azathioprine, methotrexate, cyclosporine, or oral cyclophosphamide) OR
 - B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to any standard of care drug classes listed above AND
5. ONE of the following:
 - A. Patient has NOT been treated with intravenous (IV) cyclophosphamide in the past 30 days OR
 - B. Patient has been treated with IV cyclophosphamide in the past 30 days AND will discontinue prior to starting the requested agent AND
6. ONE of the following:
 - A. Patient has NOT been treated with another biologic agent in the past 30 days OR
 - B. Patient has been treated with another biologic agent in the past 30 days AND will discontinue prior to starting the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has diagnosis of active systemic lupus erythematosus (SLE) disease AND
3. ONE of the following:
 - A. There is evidence of a claim that the patient is currently on SLE treatment regimen within the past 90 days comprised of at least ONE of the following: corticosteroids (e.g. methylprednisolone, prednisone), antimalarials (e.g. hydroxychloroquine, chloroquine), prescription nonsteroidal anti-inflammatory drugs (NSAIDS) (e.g. ibuprofen, naproxen), and/or immunosuppressives (e.g. azathioprine, methotrexate, cyclosporine, or oral cyclophosphamide) OR
 - B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to any standard of care drug classes listed above AND
4. Patient has had a decrease in symptoms or stabilization in at least ONE SLE diagnostic criteria (i.e. malar rash, discoid rash, photosensitivity, oral ulcers, nonerosive arthritis, serositis [e.g. pleuritis/pericarditis], renal disorder [e.g. persistent proteinuria greater than 0.5 grams/day or cellular casts], hematologic disorder [e.g. hemolytic anemia (with reticulocytosis), leukopenia, lymphopenia, or thrombocytopenia], and/or immunologic disorder [e.g. positive finding of antiphospholipid antibodies or anti-Sm antibodies]) AND
5. ONE of the following:
 - A. Patient has NOT been treated with intravenous (IV) cyclophosphamide in the past 30 days OR
 - B. Patient has been treated with IV cyclophosphamide in the past 30 days AND will discontinue prior to starting the requested agent AND
6. ONE of the following:
 - A. Patient has NOT been treated with another biologic agent in the past 30 days OR
 - B. Patient has been treated with another biologic agent in the past 30 days AND will discontinue prior to starting the requested agent

Prior Authorization Group – Benzodiazepines PA - Clonazepam

Drug Name(s):

**clonazepam ODT
clonazepam tablet**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently using the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

a. Seizure disorder OR

b. Panic disorder AND ONE of the following:

1) Patient's medication history includes use of a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin norepinephrine reuptake inhibitor (SNRI) OR

2) Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to formulary SSRIs or SNRIs
OR

c. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Benzodiazepines PA - Clorazepate

Drug Name(s):

clorazepate tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently using the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

a. Seizure disorder OR

b. Anxiety disorder AND ONE of the following:

1) Patient's medication history includes use of a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin norepinephrine reuptake inhibitor (SNRI) OR

2) Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to formulary SSRIs or SNRIs OR

c. Alcohol withdrawal OR

d. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Benzodiazepines PA - Diazepam

Drug Name(s):

**DIAZEPAM 1 mg/mL oral solution
diazepam intensol concentrate
diazepam tablet**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently using the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

a. Seizure disorder OR

b. Anxiety disorder AND ONE of the following:

1) Patient's medication history includes use of a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin norepinephrine reuptake inhibitor (SNRI) OR

2) Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to formulary SSRIs or SNRIs
OR

c. Skeletal muscle spasms OR

d. Alcohol withdrawal OR

e. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Benzodiazepines PA - Lorazepam

Drug Name(s):

lorazepam tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently using the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

a. Anxiety disorder AND ONE of the following:

1) Patient's medication history includes use of a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin norepinephrine reuptake inhibitor (SNRI) OR

2) Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to formulary SSRIs or SNRIs
OR

b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Benzodiazepines PA - Onfi

Drug Name(s):

clobazam suspension, tablet

ONFI

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently using the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

a. Seizure disorder OR

b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Benzodiazepines PA - Sympazan

Drug Name(s):

SYMPAZAN

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

- A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently using the requested agent AND
 - ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR
- B. BOTH of the following:
 - i. Patient has ONE of the following diagnoses:
 - a. Seizure disorder OR
 - b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Biologic Immunomodulators PA - Cosentyx

Drug Name(s):

COSENTYX

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional agent prerequisite OR
 - E. Patient's medication history indicates use of ONE formulary conventional agent prerequisite for the requested indication OR
 - F. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to at least ONE formulary conventional agent for the requested indication AND
3. ONE of the following:
 - A. Patient is NOT currently being treated with another biologic immunomodulator OR
 - B. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has shown clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. ONE of the following:

A. Patient is NOT currently being treated with another biologic immunomodulator
OR

B. Patient is currently being treated with another biologic immunomodulator AND
will discontinue the other biologic immunomodulator prior to starting the
requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE conventional agent prerequisite is required for diagnoses of psoriatic
arthritis or plaque psoriasis

NO prerequisites are required for a diagnosis of ankylosing spondylitis

Formulary conventional agents for psoriatic arthritis include cyclosporine, leflunomide,
methotrexate, or sulfasalazine

Formulary conventional topical or systemic antipsoriatic agents include acitretin,
calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Prior Authorization Group – Biologic Immunomodulators PA - Enbrel

Drug Name(s):

ENBREL

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional agent prerequisite OR
 - E. Patient's medication history indicates use of ONE formulary conventional agent prerequisite for the requested indication OR
 - F. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to at least ONE formulary conventional agent for the requested indication AND
3. Patient has been tested for latent TB AND if positive the patient has begun therapy for latent TB AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with another biologic immunomodulator OR
 - B. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has shown clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. ONE of the following:

A. Patient is NOT currently being treated with another biologic immunomodulator
OR

B. Patient is currently being treated with another biologic immunomodulator AND
will discontinue the other biologic immunomodulator prior to starting the
requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE conventional agent prerequisite is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, or juvenile idiopathic arthritis

NO prerequisites are required for a diagnosis of ankylosing spondylitis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, or psoriatic arthritis include methotrexate, hydroxychloroquine, sulfasalazine, minocycline, or leflunomide

Formulary conventional topical or systemic antipsoriatic agents include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Prior Authorization Group – Biologic Immunomodulators PA - Humira

Drug Name(s):

HUMIRA

HUMIRA KIT

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional agent prerequisite OR
 - E. Patient's medication history indicates use of ONE formulary conventional agent prerequisite for the requested indication OR
 - F. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to at least ONE formulary conventional agent for the requested indication AND
3. Patient has been tested for latent TB AND if positive the patient has begun therapy for latent TB AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with another biologic immunomodulator OR
 - B. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has shown clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. ONE of the following:

A. Patient is NOT currently being treated with another biologic immunomodulator
OR

B. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be 12 weeks for initial use for ulcerative colitis, 12 months for all others

Other Criteria:

Use of ONE conventional agent prerequisite is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, juvenile idiopathic arthritis, Crohn's disease, or ulcerative colitis

NO prerequisites are required for diagnoses of ankylosing spondylitis, hidradenitis suppurativa, or uveitis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, or psoriatic arthritis include methotrexate, hydroxychloroquine, sulfasalazine, minocycline, or leflunomide

Formulary conventional topical or systemic antipsoriatic agents include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for crohn's disease or ulcerative colitis include methotrexate

aminosalicylates, corticosteroids, cyclosporine, azathioprine, 6-mercaptopurine, metronidazole, or ciprofloxacin

Prior Authorization Group – Biologic Immunomodulators PA - Kineret

Drug Name(s):
KINERET

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND is at risk if therapy is changed OR
 - C. ONE of the following:
 - i. Patient's diagnosis is indicated in preferred biologic immunomodulator agent(s) AND ONE of the following:
 1. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR
 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to preferred biologic immunomodulator agent(s) OR
 - ii. The request is for an FDA labeled indication that is not covered by preferred biologic immunomodulator agent(s) AND
3. ONE of the following:
 - A. Patient is NOT currently being treated with another biologic immunomodulator OR
 - B. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has shown clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. ONE of the following:

A. Patient is NOT currently being treated with another biologic immunomodulator
OR

B. Patient is currently being treated with another biologic immunomodulator AND
will discontinue the other biologic immunomodulator prior to starting the
requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of TWO preferred biologics (Enbrel and Humira) is required for diagnosis of
rheumatoid arthritis

NO prerequisites are required for diagnosis of Cryopyrin-Associated Periodic
Syndromes (CAPS)/Neonatal-Onset Multisystem Inflammatory Disease (NOMID)

Prior Authorization Group – Biologic Immunomodulators PA - Orencia

Drug Name(s):

ORENCIA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND is at risk if therapy is changed OR
 - C. Patient's diagnosis is indicated in preferred biologic immunomodulator agent(s) AND ONE of the following:
 1. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR
 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to preferred biologic immunomodulator agent(s) AND
3. ONE of the following:
 - A. Patient is NOT currently being treated with another biologic immunomodulator OR
 - B. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has shown clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with another biologic immunomodulator OR

B. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of TWO preferred biologics (Enbrel and Humira) is required for the diagnoses of juvenile idiopathic arthritis or rheumatoid arthritis

Use of TWO preferred biologics (Cosentyx, Humira, Enbrel or Stelara) is required for diagnosis of psoriatic arthritis

Prior Authorization Group – Biologic Immunomodulators PA - Renflexis

Drug Name(s):

RENFLEXIS

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND is at risk if therapy is changed OR
 - C. ONE of the following:
 - i. Patient's diagnosis is indicated in preferred biologic immunomodulator agent(s) AND ONE of the following:
 1. Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR
 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to preferred biologic immunomodulator agent(s) OR
 - ii. The request is for an FDA labeled indication that is not covered by preferred biologic immunomodulator agent(s) AND
3. Patient has been tested for latent TB AND if positive the patient has begun therapy for latent TB AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with another biologic immunomodulator OR
 - B. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has shown clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

4. ONE of the following:

A. Patient is NOT currently being treated with another biologic immunomodulator
OR

B. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of TWO preferred biologics (Enbrel and Humira) is required for diagnosis of rheumatoid arthritis

Use of TWO preferred biologics (Cosentyx, Enbrel, Humira, or Stelara) is required for diagnoses of plaque psoriasis or psoriatic arthritis

Use of TWO preferred biologics (Humira and Stelara) is required for diagnosis of adult Crohn's disease

Use of TWO preferred biologics (Cosentyx, Enbrel, or Humira) is required for diagnosis of ankylosing spondylitis

Only the preferred biologic Humira is required for diagnoses of ulcerative colitis or pediatric Crohn's disease

No preferred biologic is required for the diagnosis of adult fistulizing Crohn's disease

Prior Authorization Group – Biologic Immunomodulators PA - Rituxan

Drug Name(s):

RITUXAN

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for initial approval require ONE of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

2. ALL of the following:

A. ONE of the following:

i. Patient has a diagnosis of rheumatoid arthritis AND ONE of the following:

1) Patient's medication history indicates use of preferred biologic immunomodulator agent(s) OR

2) Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to preferred biologic immunomodulator agent(s) OR

ii. Patient has another FDA labeled indication or an indication that is supported in CMS approved compendia AND

B. Patient has been screened for hepatitis B infection measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) and has begun therapy, if appropriate, prior to receiving the requested agent AND

C. ONE of the following:

i. Patient is NOT currently being treated with another biologic immunomodulator OR

ii. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent AND

D. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently being treated with the requested agent OR

C. ALL of the following:

i. Patient has been previously approved for the requested agent through the plan's PA criteria AND

ii. Patient has shown clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND

iii. ONE of the following:

a. Patient is NOT currently being treated with another biologic immunomodulator OR

b. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent AND

iv. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Use of TWO preferred biologics (Humira and Enbrel) is required for diagnosis of rheumatoid arthritis

ALL other diagnoses do NOT require any preferred biologics

Prior Authorization Group – Biologic Immunomodulators PA – Rituxan Hycela

Drug Name(s):

RITUXAN HYCELA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent OR
 - c. ALL of the following:
 - i. Patient has been screened for hepatitis B infection measuring hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (anti-HBc) and has begun therapy, if appropriate, prior to receiving the requested agent AND
 - ii. ONE of the following:
 1. Patient is NOT currently being treated with another biologic immunomodulator OR
 2. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent AND
 - iii. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent OR
 - c. ALL of the following
 - i. Patient has been previously approved for the requested agent through the plan's PA criteria AND
 - ii. Patient has shown clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
 - iii. ONE of the following:
 1. Patient is NOT currently being treated with another biologic immunomodulator OR
 2. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent AND
 - iv. Patient does NOT have any FDA labeled limitation(s) of use for the requested agent that is not otherwise supported in NCCN guidelines

There are no preferred biologics required for Rituxan Hycela

Prior Authorization Group – Biologic Immunomodulators PA - Stelara

Drug Name(s):

STELARA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's medication history indicates use of ONE formulary conventional agent prerequisite for the requested indication OR
 - E. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to at least ONE formulary conventional agent for the requested indication AND
3. ONE of the following:
 - A. Patient is NOT currently being treated with another biologic immunomodulator OR
 - B. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has shown clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with another biologic immunomodulator OR

B. Patient is currently being treated with another biologic immunomodulator AND will discontinue the other biologic immunomodulator prior to starting the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Use of ONE conventional agent prerequisite is required for diagnoses of psoriatic arthritis, plaque psoriasis, or Crohn's disease

Formulary conventional agents for psoriatic arthritis include cyclosporine, leflunomide, methotrexate, or sulfasalazine

Formulary conventional topical or systemic antipsoriatic agents include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for Crohn's disease include methotrexate, aminosalicylates, corticosteroids, cyclosporine, azathioprine, 6-mercaptopurine, metronidazole, or ciprofloxacin

Prior Authorization Group – Bivigam/Flebogamma/Gammaplex/Octagam/Privigen PA

Drug Name(s):
GAMMAPLEX

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require ONE of the following:

1. Patient has ONE of the following diagnoses:

- A. Primary immunodeficiency (including congenital agammaglobulinemia, common variable immunodeficiency (CVID), severe combined immunodeficiency, Wiskott-Aldrich Syndrome, X-linked agammaglobulinemia (XLA), humoral immunodeficiency, IgG subclass deficiency with or without IgA deficiency) OR
- B. Chronic lymphocytic leukemia OR multiple myeloma with reduced IgG AND ONE of the following:
 - i. Patient has a history of infections OR
 - ii. Patient has evidence of specific antibody deficiency OR
- C. Idiopathic thrombocytopenia purpura AND ONE of the following:
 - i. Patient has failed ONE conventional therapy (e.g. corticosteroids (e.g. methylprednisolone), or immunosuppressants (e.g. azathioprine)) OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- D. Dermatomyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy (e.g. corticosteroids (e.g. methylprednisolone) or immunosuppressants (e.g. azathioprine)) OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- E. Polymyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy (e.g. corticosteroids (e.g. methylprednisolone) or immunosuppressants (e.g. azathioprine)) OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- F. Severe rheumatoid arthritis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy (e.g. tumor necrosis factor antagonists (e.g. Humira), DMARDS (e.g. methotrexate), Renflexis) OR

- ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

Criteria continues: see Other Criteria

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 3 months for indications in NHS guidelines, 12 months all others

Other Criteria:

- G. Myasthenia gravis (MG) AND ONE of the following:
 - i. Patient is in acute myasthenic crisis with decompensation (e.g. acute episode of respiratory muscle weakness, respiratory failure, dysphagia, major functional disability responsible for the discontinuation of physical activity) OR
 - ii. Patient has severe refractory MG (e.g. major functional disability/weakness) AND ONE of the following:
 - a) Patient has failed ONE immunomodulator therapy (i.e. corticosteroid, pyridostigmine, or azathioprine) OR
 - b) Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE immunomodulator therapy OR
- H. Multiple sclerosis (MS) AND BOTH of the following:
 - i. Patient has a diagnosis of relapsing remitting MS (RRMS) AND
 - ii. Patient has had an insufficient response, documented failure, or FDA labeled contraindication(s) to TWO MS agents (e.g. Aubagio, Avonex, Betaseron, glatiramer (i.e. Copaxone, Glatopa), Plegridy, Tecfidera or Tysabri) OR
- I. Acquired von Willebrand hemophilia AND ONE of the following:
 - i. Patient has failed ONE conventional therapy (e.g. desmopressin solution, von Willebrand factor replacement therapy, corticosteroids, cyclophosphamide, FEIBA, Rituxan, or recombinant factor VIIa) OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- J. Refractory pemphigus vulgaris AND ONE of the following:
 - i. Patient has failed ONE conventional immunosuppressive therapy (e.g. azathioprine, cyclophosphamide, mycophenolate, corticosteroids) OR

ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional immunosuppressive therapy OR

2. ONE of the following:

A. Patient has another FDA labeled indication for the requested agent OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Indications with 3 month approval duration:

Idiopathic thrombocytopenia purpura, Acquired von Willebrand hemophilia, Guillain-Barre Syndrome, Lambert-Eaton myasthenia syndrome, Myasthenia gravis, Kawasaki disease, CMV induced pneumonitis in solid organ transplant, Toxic shock syndrome due to invasive group A streptococcal, Toxic epidermal necrolysis and Steven Johnsons syndrome

Drug is also subject to Part B versus Part D review.

Prior Authorization Group – Carbaglu PA

Drug Name(s):

CARBAGLU

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of ONE of the following:
 - a. Acute hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS) OR
 - b. Chronic hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS) AND
2. The dose requested is within the FDA labeled dosing for the requested indication

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. geneticist, metabolic disorders) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Chenodal PA

Drug Name(s):

CHENODAL

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of radiolucent stones in a well-opacifying gallbladder AND
2. The dose requested is within FDA labeled dosing

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Chorionic Gonadotropin PA

Drug Name(s):

**CHORIONIC GONADOTROPIN
PREGNYL**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Used to promote fertility or to treat erectile dysfunction AND FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ONE of the following:

A. Patient has a diagnosis of prepubertal cryptorchidism not due to anatomic obstruction
OR

B. Patient is a male with a diagnosis of hypogonadotropic hypogonadism (hypogonadism secondary to pituitary deficiency) AND BOTH of the following:

i. Patient has a measured current or pretreatment total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR a free serum testosterone level that is below the testing laboratory's lower limit of the normal range AND

ii. Patient has measured luteinizing hormone (LH) AND follicle-stimulating hormone (FSH) levels that are at (low-normal) or below the testing laboratory's normal range OR

C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Cinacalcet PA

Drug Name(s):

**cinacalcet tablet
SENSIPAR**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require the following:

1. Patient has ONE of the following:

A. An FDA approved indication or has an indication that is supported in CMS approved compendia for the requested agent not otherwise excluded from Part D [i.e., secondary hyperparathyroidism due to end-stage renal disease (ESRD) on dialysis] AND ONE of the following:

i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

ii. Prescriber states the patient is currently using the requested agent AND is at risk if therapy is changed OR

B. A diagnosis of hypercalcemia due to parathyroid carcinoma OR

C. A diagnosis of primary hyperparathyroidism (HPT) AND BOTH of the following:

i. Patient has a pretreatment serum calcium that is above the testing laboratory's upper limit of normal AND

ii. Patient is unable to undergo parathyroidectomy OR

D. Another indication that is supported in CMS approved compendia for the requested agent not otherwise excluded from Part D

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Corlanor PA

Drug Name(s):

CORLANOR

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. Patient has stable, symptomatic chronic heart failure (e.g. NYHA Class II, III, IV, ACCF/AHA Class C, D) AND
2. ONE of the following:
 - a. ALL of the following:
 - i. The requested agent is for a pediatric patient, 6 months or over AND
 - ii. Patient has heart failure due to dilated cardiomyopathy (DCM) AND
 - iii. Patient is in sinus rhythm with an elevated heart rate OR
 - b. ALL of the following:
 - i. The requested agent is for an adult patient AND
 - ii. Patient has a baseline OR current left ventricular ejection fraction of 35% or less AND
 - iii. Patient is in sinus rhythm with a resting heart rate of 70 beats or greater per minute prior to initiating therapy with the requested agent AND
 - iv. ONE of the following:
 1. Patient is on a maximally tolerated dose of beta blocker (e.g. atenolol, bisoprolol, carvedilol, metoprolol) OR
 2. Patient has a history of a documented intolerance, FDA labeled contraindication, or hypersensitivity to a beta blocker

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Cresemba PA

Drug Name(s):

CRESEMBA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ONE of the following:

1. Patient has a diagnosis of invasive aspergillosis OR
2. Patient has a diagnosis of invasive mucormycosis OR
3. Patient has another indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. ONE of the following:
 - A. Patient has a diagnosis of invasive aspergillosis and the patient has continued indicators of active disease (e.g. continued radiologic findings, direct microscopy findings, histopathology findings, positive cultures, positive serum galactomannan assay) OR
 - B. Patient has a diagnosis of invasive mucormycosis and the patient has continued indicators of active disease (e.g. continued radiologic findings, direct microscopy findings, histopathology findings, positive cultures, positive serum galactomannan assay) OR
 - C. Patient has another indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 6 months

Other Criteria:

Prior Authorization Group – Crysvita PA

Drug Name(s):

CRYSVITA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of X-linked hypophosphatemia (XLH) as confirmed by testing for renal phosphate wasting AND radiography AND
2. ONE of the following:
 - a. Patient's epiphyseal plate has not fused OR
 - b. Patient's epiphyseal plate has fused AND the patient is experiencing symptoms of XLH (e.g. bone pain, fractures, limited mobility) AND
3. The dose requested is within the FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's prior authorization criteria AND
2. Patient has a diagnosis of X-linked hypophosphatemia (XLH) AND
3. Patient has had clinical improvement with the requested agent (e.g. enhanced height velocity, improvement in lower extremity bowing and associated abnormalities, radiographic evidence of epiphyseal healing, improvement in bone pain, enhanced mobility, improvement in osteomalacia, and improvement in fracture healing) AND
4. The dose requested is within the FDA labeled dosing for the requested indication

Age Restrictions:

Patient is 1 year of age or older

Prescriber Restrictions:

Prescriber is a specialist (e.g. nephrologist, endocrinologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Cystinosis Agents PA - Cystagon

Drug Name(s):

CYSTAGON

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of nephropathic cystinosis AND
2. Prescriber has performed a baseline white blood cell (WBC) cystine level test AND
3. The dose requested is within the FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of nephropathic cystinosis AND
3. Patient has had clinical improvement (e.g. decrease in WBC cystine levels from baseline) with the requested agent AND
4. The dose requested is within the FDA labeled dosing for the requested indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Daklinza PA

Drug Name(s):

DAKLINZA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of chronic hepatitis C infection confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The dosing of the requested agent is within the FDA labeled dosing or is AASLD/IDSA guideline supported AND
5. ONE of the following:
 - A. There is documentation that the patient is currently being treated with the requested agent OR
 - B. Patient has an FDA labeled contraindication or hypersensitivity to the preferred agents: Harvoni (ledipasvir/sofosbuvir) for supported genotypes OR Epclusa (sofosbuvir/velpatasvir) for genotype 2 or 3 OR
 - C. Prescriber has submitted documentation based on FDA approved labeling or AASLD/IDSA guidelines to support the use of the non-preferred agent for the patient's diagnosis and genotype over the preferred agents: Harvoni (ledipasvir/sofosbuvir) for supported genotypes OR Epclusa (sofosbuvir/velpatasvir) for genotype 2 or 3

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group – Dalfampridine PA

Drug Name(s):

AMPYRA

dalfampridine ER tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require BOTH of the following:

1. Patient has a diagnosis of multiple sclerosis (MS) AND
2. If the patient has relapsing form of MS, ONE of the following:
 - A. There is evidence of a claim that the patient is receiving concurrent therapy within the past 30 days with a disease modifying agent (e.g. Aubagio, Avonex, Betaseron, Extavia, Gilenya, glatiramer (i.e. Copaxone, Glatopa), Lemtrada, mitoxantrone, Plegridy, Rebif, Tecfidera, or Tysabri) OR
 - B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a disease modifying agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of multiple sclerosis (MS) AND
3. Patient has demonstrated a stabilization or improvement from baseline in timed walking speed (timed 25 foot walk)

Age Restrictions:

Prescriber Restrictions:

Prescriber is a neurologist or the prescriber has consulted with a neurologist

Coverage Duration:

Initial approval 3 months. 12 months for renewal.

Other Criteria:

Prior Authorization Group – Daliresp PA

Drug Name(s):

DALIRESP

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of following:
 - a. Patient has had an inadequate response to an agent from two of the following categories:
 - i. long-acting beta-2 agonist (LABA) [e.g. salmeterol]
 - ii. long-acting antimuscarinic antagonist/anticholinergic (LAMA) [e.g. tiotropium]
 - iii. inhaled corticosteroid (ICS) [e.g. fluticasone] OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to an agent from two of the following categories:
 - i. long-acting beta-2 agonist (LABA) [e.g. salmeterol]
 - ii. long-acting antimuscarinic antagonist/anticholinergic (LAMA) [e.g. tiotropium]
 - iii. inhaled corticosteroid (ICS) [e.g. fluticasone]

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Dupixent PA

Drug Name(s):

DUPIXENT

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require BOTH of the following:

1. ONE of the following:

A. Patient has a diagnosis of moderate-to-severe atopic dermatitis AND BOTH of the following:

i. ONE of the following:

a. Patient has tried and failed a topical steroid (e.g. clobetasol, triamcinolone) OR

b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a topical steroid AND

ii. ONE of the following:

a. Patient has tried and failed a topical calcineurin inhibitor (e.g. pimecrolimus) OR

b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a topical calcineurin inhibitor OR

Criteria continues: see Other Criteria

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. dermatologist, allergist, immunologist, pulmonologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

B. Patient has a diagnosis of moderate-to-severe asthma AND ALL of the following:

i. Patient is 12 years of age or over AND

ii. ONE of the following:

- a. Patient has an eosinophilic phenotype AND the patient has a baseline blood eosinophil count of 150 cells/microliter or higher OR
 - b. Patient has oral corticosteroid dependent asthma AND
 - iii. Patient has a baseline Forced Expiratory Volume (FEV1) that is less than 80% of predicted AND
 - iv. ONE of the following:
 - a. BOTH of the following:
 - 1. Patient is NOT currently being treated with the requested agent AND
 - 2. Patient is currently being treated with a maximally tolerated inhaled corticosteroid within the past 90 days OR
 - b. BOTH of the following:
 - 1. Patient is currently being treated with the requested agent AND
 - 2. Patient is also currently being treated with an inhaled corticosteroid that is dosed as needed to control symptoms OR
 - c. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to an inhaled corticosteroid AND
 - v. ONE of the following:
 - a. There is evidence of a claim within the past 90 days that the patient is currently being treated with ONE of the following:
 - 1. A long-acting beta-2 agonist (LABA) OR
 - 2. A leukotriene receptor antagonist (LRTA) OR
 - 3. A long-acting muscarinic antagonist (LAMA) OR
 - 4. Theophylline OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a long-acting beta-2 agonist (LABA), leukotriene receptor antagonist (LRTA), long-acting muscarinic antagonist (LAMA), OR theophylline AND
 - vi. The requested agent will NOT be used in combination with Xolair or with an injectable interleukin 5 (IL-5) inhibitor (e.g. Cinqair, Fasenra, Nucala) for the requested indication AND
- 2. The dose requested is within the FDA labeled dosing for the requested indication

Prior Authorization Group – Egrifta PA

Drug Name(s):

EGRIFTA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of HIV infection AND
2. The requested agent will be used to reduce excess abdominal fat in an HIV-infected patient with lipodystrophy AND
3. Patient is currently on anti-retroviral therapy (ART) within the past 90 days AND
4. Prescriber has measured baseline visceral adipose tissue (VAT) and waist circumference AND
5. Patient is NOT planning to become pregnant or currently breastfeeding

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. The requested agent will be used to reduce excess abdominal fat in an HIV-infected patient with lipodystrophy AND
3. Patient is currently on anti-retroviral therapy (ART) within the past 90 days AND
4. Patient has achieved or maintained a decrease in visceral adipose tissue (VAT) from baseline OR maintained or decreased waist circumference from baseline AND
5. Patient is NOT planning to become pregnant or currently breastfeeding

Age Restrictions:

Patient is between 18 and 65 years of age

Prescriber Restrictions:

Coverage Duration:

Initial approval 6 months. Renewal approval 12 months.

Other Criteria:

Prior Authorization Group – Emgality PA

Drug Name(s):

EMGALITY

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of migraine AND
2. Patient has 4 migraine headaches or more per month AND
3. ONE of the following:
 - a. Patient has failed at least TWO conventional migraine prophylaxis agents from TWO different classes (e.g. beta blockers [propranolol], anticonvulsants [divalproex, topiramate]) OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to TWO conventional migraine prophylaxis agents from TWO different classes AND
4. ONE of the following:
 - a. Patient is NOT currently taking another calcitonin gene-related peptide (CGRP) agent OR
 - b. Patient is currently being treated with another calcitonin gene-related peptide (CGRP) agent AND will discontinue prior to initiating therapy with the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's prior authorization criteria AND
2. Patient has a diagnosis of migraine AND
3. Patient has shown clinical benefit with the requested agent AND
4. ONE of the following:
 - a. Patient is NOT currently taking another calcitonin gene-related peptide (CGRP) agent OR
 - b. Patient is currently being treated with another calcitonin gene-related peptide (CGRP) agent AND will discontinue prior to continuing therapy with the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Epclusa PA

Drug Name(s):

EPCLUSA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The dosing of the requested agent is within the FDA labeled dosing or is AASLD/IDSA guideline supported AND
5. ONE of the following:
 - A. There is documentation that the patient is currently being treated with the requested agent OR
 - B. Patient has an FDA labeled contraindication or hypersensitivity to the preferred agent: Harvoni (ledipasvir/sofosbuvir) for supported genotypes OR
 - C. Prescriber has submitted documentation based on FDA approved labeling or AASLD/IDSA guidelines to support the use of the non-preferred agent for the patient's diagnosis and genotype over the preferred agent: Harvoni (ledipasvir/sofosbuvir) for supported genotypes

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group – Epidiolex PA

Drug Name(s):

EPIDIOLEX

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.

Criteria for approval require BOTH of the following:

1. ONE of the following:
 - a. Patient has a diagnosis of Lennox-Gastaut syndrome OR
 - b. Patient has a diagnosis of Dravet syndrome AND
2. The dose requested is within the FDA labeled dosing for the requested indication

Age Restrictions:

Patient is 2 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Erythropoietin Stimulating Agents PA - Aranesp

Drug Name(s):

ARANESP

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. The requested agent is being prescribed for ONE of the following:

A. Anemia due to myelosuppressive chemotherapy for a non-myeloid malignancy
AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR less than 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

ii. Patient is being concurrently treated with chemotherapy, with or without radiation (treatment period extends to 8 weeks post chemotherapy) AND

iii. Intent of chemotherapy is non-curative OR

B. Anemia associated with chronic renal failure in a patient NOT on dialysis AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR 11 g/dL or less for patients stabilized on therapy (measured within the previous 4 weeks) AND

ii. Rate of hemoglobin decline indicates the likelihood of requiring a RBC transfusion AND

iii. Intent of therapy is to reduce risk of alloimmunization and/or other RBC transfusion related risks OR

C. Anemia due to myelodysplastic syndrome AND patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks)
OR

D. Another indication that is supported in CMS approved compendia for the requested agent AND patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

2. Patient's transferrin saturation and serum ferritin have been evaluated

Drug is also subject to Part B versus Part D review.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

6 months for chemotherapy, 12 months for other indications

Other Criteria:

Prior Authorization Group – Erythropoietin Stimulating Agents PA-Epogen/Procrit

Drug Name(s):

EPOGEN

PROCRIT

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of:

1. Requested agent is being prescribed for ONE of:

A. To reduce the possibility of allogeneic blood transfusion in surgery patient (pt) AND pt's hemoglobin level is greater than 10 g/dL but 13 g/dL or less OR

B. Anemia due to myelosuppressive chemotherapy for a non-myeloid malignancy AND ALL of:

i. Pt's hemoglobin level is less than 10 g/dL for pts initiating ESA therapy OR less than 12 g/dL for pts stabilized on therapy (measured within the previous 4 weeks) AND

ii. Pt is being concurrently treated with chemotherapy with or without radiation (treatment period extends to 8 weeks post chemotherapy) AND

iii. Intent of chemotherapy is non-curative OR

C. Anemia associated with chronic renal failure in a pt NOT on dialysis AND ALL of:

i. Pt's hemoglobin level is less than 10 g/dL for pts initiating ESA therapy OR 11 g/dL or less for pts stabilized on therapy (measured within the previous 4 weeks) AND

ii. Rate of hemoglobin decline indicates the likelihood of requiring a RBC transfusion AND

iii. Intent of therapy is to reduce the risk of alloimmunization and/or other RBC transfusion related risks OR

D. Anemia due to myelodysplastic syndrome AND pt's hemoglobin level is less than 12 g/dL for pts initiating ESA therapy OR less than or equal to 12 g/dL for pts stabilized on therapy (measured within previous 4 weeks) OR

E. Anemia resulting from zidovudine treatment of HIV infection AND pt's hemoglobin level is less than 12 g/dL for pts initiating ESA therapy OR less than or equal to 12 g/dL for pts stabilized on therapy (measured within previous 4 weeks) OR

F. Another indication supported in CMS approved compendia for requested agent AND pt's hemoglobin level is less than 12 g/dL for pts initiating ESA therapy OR less than or equal to 12 g/dL for pts stabilized on therapy (measured within previous 4 weeks) AND

2. Pt's transferrin saturation and serum ferritin have been evaluated

Drug is also subject to Part B versus Part D review.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

1 month for surgery (reduce transfusion possibility), 6 months for chemo, 12 months for other

Other Criteria:

Prior Authorization Group – Erythropoietin Stimulating Agents PA - Retacrit

Drug Name(s):

RETACRIT

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. The requested agent is being prescribed for ONE of the following:

A. To reduce the possibility of allogeneic blood transfusion in a surgery patient AND the patient's hemoglobin level is greater than 10 g/dL but 13 g/dL or less OR

B. Anemia due to myelosuppressive chemotherapy for a non-myeloid malignancy AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR less than 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

ii. Patient is being concurrently treated with chemotherapy with or without radiation (treatment period extends to 8 weeks post chemotherapy) AND

iii. The intent of chemotherapy is non-curative OR

C. Anemia associated with chronic renal failure in a patient NOT on dialysis AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR 11 g/dL or less for patients stabilized on therapy (measured within the previous 4 weeks) AND

ii. The rate of hemoglobin decline indicates the likelihood of requiring a RBC transfusion AND

iii. The intent of therapy is to reduce the risk of alloimmunization and/or other RBC transfusion related risks OR

D. Anemia resulting from zidovudine treatment of HIV infection AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR 12 g/dL or less for patients stabilized on therapy (measured within the previous 4 weeks) OR

E. Another indication that is supported in CMS approved compendia for the requested agent AND the patient's hemoglobin level is less than 12 g/dL for

patients initiating ESA therapy OR 12 g/dL or less for patients stabilized on therapy (measured within the previous 4 weeks) AND
2. Patient's transferrin saturation and serum ferritin have been evaluated

Drug is also subject to Part B versus Part D review.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

1 month for surgery (reduce transfusion possibility), 6 months for chemo, 12 months for other

Other Criteria:

Prior Authorization Group – Fentanyl Nasal PA - Lazanda

Drug Name(s):

LAZANDA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require BOTH of the following:

1. ONE of the following:

- a. Patient has a diagnosis of chronic cancer pain due to an active malignancy AND there is evidence of a claim that the patient is currently taking a long-acting opioid with the nasal fentanyl within the past 90 days OR
- b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- a. Patient is NOT currently receiving any other oral or nasal fentanyl agent OR
- b. Patient is currently receiving another oral or nasal fentanyl agent AND will be discontinued prior to initiating the requested agent

Age Restrictions:

Patient is 18 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Fentanyl Oral PA - Abstral

Drug Name(s):

ABSTRAL

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require BOTH of the following:

1. ONE of the following:

- a. Patient has a diagnosis of chronic cancer pain due to an active malignancy AND there is evidence of a claim that the patient is currently taking a long-acting opioid with the oral fentanyl within the past 90 days OR
- b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- a. Patient is NOT currently receiving any other oral or nasal fentanyl agent OR
- b. Patient is currently receiving another oral or nasal fentanyl agent AND will be discontinued prior to initiating the requested agent

Age Restrictions:

Patient is 18 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Fentanyl Oral PA - Fentanyl lozenge

Drug Name(s):

fentanyl citrate oral lozenge

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require BOTH of the following:

1. ONE of the following:

- a. Patient has a diagnosis of chronic cancer pain due to an active malignancy AND there is evidence of a claim that the patient is currently taking a long-acting opioid with the oral fentanyl within the past 90 days OR
- b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- a. Patient is NOT currently receiving any other oral or nasal fentanyl agent OR
- b. Patient is currently receiving another oral or nasal fentanyl agent AND will be discontinued prior to initiating the requested agent

Age Restrictions:

Patient is 16 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Forteo PA

Drug Name(s):

FORTEO

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

An increased baseline risk for osteosarcoma

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:
 - a. Postmenopausal female with osteoporosis OR
 - b. Male with primary or hypogonadal osteoporosis OR
 - c. Osteoporosis with sustained systemic glucocorticoid therapy AND
2. ONE of the following:
 - a. Patient has a history of vertebral fracture(s), or low trauma or fragility fracture(s) [e.g. prior fracture from minor trauma such as falling from standing height or less] within the past 5 years OR
 - b. Patient has a T-score that is -2.5 or lower (greater than or equal to 2.5 SD below the mean BMD value for a young adult) AND ONE of the following:
 - i. Patient is female and has failed either a bisphosphonate or SERM OR
 - ii. Patient is male and has failed a bisphosphonate OR
 - iii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a SERM or bisphosphonate (bisphosphonate or SERM for female patients, bisphosphonates only for male patients) AND
3. ONE of the following:
 - a. Patient is not receiving concomitant bisphosphonate, SERM, Prolia (denosumab), or Xgeva (denosumab) therapy within the past 90 days OR
 - b. Prescriber indicates that the patient will discontinue the current bisphosphonate, SERM, Prolia (denosumab) or Xgeva (denosumab) therapy prior to initiating therapy with the requested agent AND
4. The dose requested is within the FDA approved labeling AND
5. The total duration of treatment with Forteo has not exceeded 2 years

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

No prior Forteo use approve 2 yrs, Prior Forteo use approve remainder of 2 yrs of total therapy

Other Criteria:

Prior Authorization Group – Gammagard/Gammaked/Gamunex-C PA

Drug Name(s):

GAMMAGARD
GAMMAGARD SD
GAMUNEX-C

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require ONE of the following:

1. Patient has ONE of the following diagnoses:

- A. Primary immunodeficiency (including congenital agammaglobulinemia, common variable immunodeficiency (CVID), severe combined immunodeficiency, Wiskott-Aldrich Syndrome, X-linked agammaglobulinemia (XLA), humoral immunodeficiency, IgG subclass deficiency with or without IgA deficiency) OR
- B. Chronic lymphocytic leukemia OR multiple myeloma with reduced IgG AND ONE of the following:
 - i. Patient has a history of infections OR
 - ii. Patient has evidence of specific antibody deficiency OR
- C. Idiopathic thrombocytopenia purpura AND ONE of the following:
 - i. Patient has failed ONE conventional therapy (e.g. corticosteroids (e.g. methylprednisolone), or immunosuppressants (e.g. azathioprine)) OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- D. Dermatomyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy (e.g. corticosteroids (e.g. methylprednisolone) or immunosuppressants (e.g. azathioprine)) OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- E. Polymyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy (e.g. corticosteroids (e.g. methylprednisolone) or immunosuppressants (e.g. azathioprine)) OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- F. Severe rheumatoid arthritis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy (e.g. tumor necrosis factor antagonists (e.g. Humira), DMARDS (e.g. methotrexate), Renflexis) OR

- ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

Criteria continues: see Other Criteria

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 3 months for indications in NHS guidelines, 12 months all others

Other Criteria:

- G. Myasthenia gravis (MG) AND ONE of the following:
 - i. Patient is in acute myasthenic crisis with decompensation (e.g. acute episode of respiratory muscle weakness, respiratory failure, dysphagia, major functional disability responsible for the discontinuation of physical activity) OR
 - ii. Patient has severe refractory MG (e.g. major functional disability/weakness) AND ONE of the following:
 - a) Patient has failed ONE immunomodulator therapy (i.e. corticosteroid, pyridostigmine, or azathioprine) OR
 - b) Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE immunomodulator therapy OR
 - H. Multiple sclerosis (MS) AND BOTH of the following:
 - i. Patient has a diagnosis of relapsing remitting MS (RRMS) AND
 - ii. Patient has had an insufficient response, documented failure, or FDA labeled contraindication(s) to TWO MS agents (e.g. Aubagio, Avonex, Betaseron, glatiramer (i.e. Copaxone, Glatopa), Plegridy, Tecfidera or Tysabri) OR
 - I. Acquired von Willebrand hemophilia AND ONE of the following:
 - i. Patient has failed ONE conventional therapy (e.g. desmopressin solution, von Willebrand factor replacement therapy, corticosteroids, cyclophosphamide, FEIBA, Rituxan, or recombinant factor VIIa) OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
 - J. Refractory pemphigus vulgaris AND ONE of the following:
 - i. Patient has failed ONE conventional immunosuppressive therapy (e.g. azathioprine, cyclophosphamide, mycophenolate, corticosteroids) OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional immunosuppressive therapy OR
2. ONE of the following:
- A. Patient has another FDA labeled indication for the requested agent OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Indications with 3 month approval duration:

Idiopathic thrombocytopenia purpura, Acquired von Willebrand hemophilia, Guillain-Barre Syndrome, Lambert-Eaton myasthenia syndrome, Myasthenia gravis, Kawasaki disease, CMV induced pneumonitis in solid organ transplant, Toxic shock syndrome due to invasive group A streptococcal, Toxic epidermal necrolysis and Steven Johnsons syndrome

Drug is also subject to Part B versus Part D review.

Prior Authorization Group – Gattex PA

Drug Name(s):

GATTEX

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of short bowel syndrome (SBS) AND
2. Patient is dependent on parenteral nutrition OR intravenous (PN/IV) fluids AND
3. Patient has had a colonoscopy with any polyps removed, if present, within the last 6 months

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of short bowel syndrome (SBS) AND
3. Patient has had a reduction from baseline in parenteral nutrition OR intravenous (PN/IV) fluids

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Initial approval 6 months. Renewal approval 12 months.

Other Criteria:

Prior Authorization Group – Gaucher Enzyme Replacement PA - Cerezyme

Drug Name(s):

CEREZYME

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of Gaucher disease type 1 confirmed by ONE of the following:
 - A. A baseline glucocerebrosidase enzyme activity of less than 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR
 - B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND
2. Prescriber has drawn baseline levels of hemoglobin, platelets, liver volume, and spleen volume AND
3. Prior to any treatment for the intended diagnosis, the patient has had ONE of the following:
 - A. Anemia (defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender) OR
 - B. Thrombocytopenia (defined as a platelet count of less than 100,000 per microliter) OR
 - C. Hepatomegaly OR
 - D. Splenomegaly OR
 - E. Growth failure (growth velocity below the standard mean for age) OR
 - F. Evidence of bone disease with other causes ruled out

Criteria for renewal require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of Gaucher's disease type 1 AND
3. Patient has shown clinical improvement and/or stabilization from baseline in ONE of the following:
 - A. Hemoglobin (Hb) levels OR
 - B. Platelet count OR
 - C. Hepatomegaly OR
 - D. Splenomegaly OR
 - E. Growth OR

F. Bone pain or disease

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. endocrinologist, geneticist, hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Gaucher Enzyme Replacement PA - Elelyso

Drug Name(s):

ELELYSO

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of Gaucher disease type 1 confirmed by ONE of the following:
 - A. A baseline glucocerebrosidase enzyme activity of less than 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR
 - B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND
2. Prescriber has drawn baseline levels of hemoglobin, platelets, liver volume, and spleen volume AND
3. Prior to any treatment for the intended diagnosis, the patient has had ONE of the following:
 - A. Anemia (defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender) OR
 - B. Thrombocytopenia (defined as a platelet count of less than 100,000 per microliter) OR
 - C. Hepatomegaly OR
 - D. Splenomegaly OR
 - E. Growth failure (growth velocity below the standard mean for age) OR
 - F. Evidence of bone disease with other causes ruled out

Criteria for renewal require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of Gaucher's disease type 1 AND
3. Patient has shown clinical improvement and/or stabilization from baseline in ONE of the following:
 - A. Hemoglobin (Hb) levels OR
 - B. Platelet count OR
 - C. Hepatomegaly OR
 - D. Splenomegaly OR
 - E. Growth OR

F. Bone pain or disease

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. endocrinologist, geneticist, hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Gaucher Enzyme Replacement PA - Vpriv

Drug Name(s):

VPRIV

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of Gaucher disease type 1 confirmed by ONE of the following:
 - A. A baseline glucocerebrosidase enzyme activity of less than 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR
 - B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND
2. Prescriber has drawn baseline levels of hemoglobin, platelets, liver volume, and spleen volume AND
3. Prior to any treatment for the intended diagnosis, the patient has had ONE of the following:
 - A. Anemia (defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender) OR
 - B. Thrombocytopenia (defined as a platelet count of less than 100,000 per microliter) OR
 - C. Hepatomegaly OR
 - D. Splenomegaly OR
 - E. Growth failure (growth velocity below the standard mean for age) OR
 - F. Evidence of bone disease with other causes ruled out

Criteria for renewal require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of Gaucher's disease type 1 AND
3. Patient has shown clinical improvement and/or stabilization from baseline in ONE of the following:
 - A. Hemoglobin (Hb) levels OR
 - B. Platelet count OR
 - C. Hepatomegaly OR
 - D. Splenomegaly OR
 - E. Growth OR

F. Bone pain or disease

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. endocrinologist, geneticist, hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Growth Hormone PA - Omnitrope

Drug Name(s):

OMNITROPE

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

For Children – Criteria for initial approval require the following:

1. ONE of the following:

- a. Patient is a neonate (4 months of age or younger) with hypoglycemia in the absence of metabolic disorder AND
 - i. The GH level is less than 20 ng/mL OR
- b. Patient has a diagnosis of Turner Syndrome OR
- c. Patient has a diagnosis of Prader-Willi Syndrome OR
- d. Patient has a diagnosis of panhypopituitarism AND BOTH of the following:
 - i. Deficiencies in 3 or more pituitary axes AND
 - ii. Measured serum IGF-1 (insulin-like growth factor-1) levels are below the age and sex-appropriate reference range when off GH therapy OR
- e. Patient has a diagnosis of growth hormone deficiency (GHD) or short stature AND BOTH of the following:
 - i. Patient has ONE of the following:
 - a) Height more than 2 standard deviations (SD) below the mean for age and sex OR
 - b) Height more than 1.5 SD below the midparental height OR
 - c) A decrease in height SD of more than 0.5 over one year in children at least 2 years of age OR
 - d) Height velocity more than 2 SD below the mean over one year or more than 1.5 SD sustained over two years AND
 - ii. Failure of at least 2 growth hormone (GH) stimulation tests (peak GH value of less than 10 mcg/L after stimulation) OR
- f. Patient has a diagnosis of small for gestational age (SGA) AND ALL of the following:
 - i. Patient is at least 2 years of age AND
 - ii. Documented birth weight and/or length that is 2 or more standard deviations (SD) below the mean for gestational age AND

- iii. At 24 months of age, the patient fails to manifest catch-up growth evidenced by a height 2 or more SD below the mean for age and sex

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

For Children – Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the preferred agent through the plan's PA criteria AND
2. Patient has been diagnosed with ONE of the following:
 - a. Neonate (4 months of age or younger) with hypoglycemia in the absence of metabolic disorder OR
 - b. Growth Hormone Deficiency, Short Stature OR
 - c. Panhypopituitarism OR
 - d. Prader-Willi Syndrome OR
 - e. Small for Gestational Age (SGA) OR
 - f. Turner Syndrome AND
3. ALL of the following:
 - a. Patient does not have closed epiphyses AND
 - b. Patient is being monitored for adverse effects of therapy with the requested agent AND
 - c. Patient's height is increased or height velocity has improved since initiation or last approval of the requested agent

For Adults – Criteria for initial approval require the following:

1. Patient has been diagnosed with ONE of the following:
 - a. Childhood growth hormone deficiency (GHD) with genetic or organic origin AND ONE of the following:
 - i. Low IGF-1 (insulin-like growth factor-1) level without GH replacement therapy OR
 - ii. Failure of at least one growth hormone (GH) stimulation test as an adult (peak GH value of 5 mcg/L or lower after stimulation) OR
 - b. Acquired adult GHD secondary to structural lesions or trauma AND ONE of the following:
 - i. Patient has a diagnosis of panhypopituitarism AND BOTH of the following:

- a) Deficiencies in 3 or more pituitary axes AND
- b) Low IGF-1 level without GH replacement therapy OR
- ii. Patient has failed at least one growth hormone (GH) stimulation test as an adult OR
- c. Idiopathic GHD (adult or childhood onset) AND the patient has failed at least two growth hormone (GH) stimulation tests as an adult

For Adults – Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the preferred agent through the plan's PA criteria AND
2. Patient has been diagnosed with ONE of the following:
 - a. Childhood growth hormone deficiency (GHD) with genetic or organic origin OR
 - b. Acquired adult GHD secondary to structural lesions or trauma OR
 - c. Other (e.g., childhood idiopathic GHD, adult-onset idiopathic GHD) AND
3. Patient is being monitored for adverse effects of therapy with the requested agent AND
4. Patient's IGF-1 level has been evaluated to confirm the appropriateness of the current dose AND
5. Patient has had benefits from therapy with the requested agent in any of the following response parameters: body composition, hip-to-waist ratio, cardiovascular health, bone mineral density, serum cholesterol, physical strength, or quality of life

Prior Authorization Group – HAE PA - Cinryze

Drug Name(s):

CINRYZE

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D, plus the off-label use for acute HAE attacks.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of hereditary angioedema (HAE) which has been diagnosed with measurement of C1-INH antigenic level, C1-INH functional level, and C4 level as follows:
 - a. Type I HAE: Two separate measurements indicating decreased quantities of C4 and C1-INH OR
 - b. Type II HAE: Two separate measurements indicating decreased quantities of C4 and decreased C1-INH function (C1-INH protein level may be normal) OR
 - c. Type III HAE: Normal levels of C4 and C1-INH (at baseline and during an attack) AND ONE of the following:
 - i. Family history of angioedema OR
 - ii. Patient demonstrates a Factor XII mutation that is associated with the disease AND
2. Medications known to cause angioedema (i.e. ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued when appropriate AND
3. ONE of the following:
 - a. The requested agent will be used to treat acute HAE attacks AND ONE of the following:
 - i. Patient is receiving only ONE agent indicated for treatment of acute HAE OR
 - ii. The other agent being used for acute HAE attacks will be discontinued before starting the requested agent OR
 - b. The requested agent will be used for prophylaxis against HAE attacks AND ALL of the following:
 - i. ONE of the following:
 1. Patient is receiving only ONE HAE agent indicated for prophylaxis against HAE attacks OR
 2. The other agent being used for prophylaxis will be discontinued before starting the requested agent AND

- ii. Patient has a history of at least 2 acute attacks per month OR 2 severe attacks (laryngeal or abdominal attacks) per year OR for short-term prophylaxis if prior to medical, surgical, or dental procedure AND
- iii. ONE of the following:
 - 1. Patient has tried a formulary 17 alpha-alkylated androgen or antifibrinolytic agent OR
 - 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a formulary 17 alpha-alkylated androgen or antifibrinolytic agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
- 2. Patient has a diagnosis of hereditary angioedema (HAE) AND ONE of the following:
 - a. The requested indication is for acute HAE AND ONE of the following:
 - i. Patient is receiving only ONE agent indicated for treatment of acute HAE OR
 - ii. The other agent being used for acute HAE attacks will be discontinued before starting the requested agent OR
 - b. The requested indication is for prophylaxis of HAE attacks AND ONE of the following:
 - i. Patient is receiving only ONE HAE agent indicated for prophylaxis against HAE attacks OR
 - ii. The other agent being used for prophylaxis will be discontinued before starting the requested agent AND
- 3. Patient has had a decrease in the frequency of acute attacks or has had stabilization of disease from use of the requested agent

Prior Authorization Group – HAE PA - Firazyr

Drug Name(s):

FIRAZYR

icatibant 30 mg/3 mL injection

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of hereditary angioedema (HAE) which has been diagnosed with measurement of C1-INH antigenic level, C1-INH functional level, and C4 level as follows:
 - a. Type I HAE: Two separate measurements indicating decreased quantities of C4 and C1-INH OR
 - b. Type II HAE: Two separate measurements indicating decreased quantities of C4 and decreased C1-INH function (C1-INH protein level may be normal) OR
 - c. Type III HAE: Normal levels of C4 and C1-INH (at baseline and during an attack) AND ONE of the following:
 - i. Family history of angioedema OR
 - ii. Patient demonstrates a Factor XII mutation that is associated with the disease AND
2. Medications known to cause angioedema (i.e. ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued when appropriate AND
3. The requested agent will be used to treat acute HAE attacks AND
4. ONE of the following:
 - a. Patient is receiving only ONE agent indicated for treatment of acute HAE attacks OR
 - b. The other agent being used for acute HAE attacks will be discontinued before starting the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of hereditary angioedema (HAE) AND
3. The requested agent will be used to treat acute HAE attacks AND
4. ONE of the following:
 - a. Patient is receiving only ONE agent indicated for treatment of acute HAE OR
 - b. The other agent being used for acute HAE attacks will be discontinued before starting the requested agent AND
5. Patient has had a decrease in the frequency of acute attacks or stabilization of disease from use of the requested agent

Prior Authorization Group – HAE PA - Haegarda

Drug Name(s):

HAEGARDA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of hereditary angioedema (HAE) which has been diagnosed with measurement of C1-INH antigenic level, C1-INH functional level, and C4 level as follows:
 - a. Type I HAE: Two separate measurements indicating decreased quantities of C4 and C1-INH OR
 - b. Type II HAE: Two separate measurements indicating decreased quantities of C4 and decreased C1-INH function (C1-INH protein level may be normal) OR
 - c. Type III HAE: Normal levels of C4 and C1-INH (at baseline and during an attack) AND ONE of the following:
 - i. Family history of angioedema OR
 - ii. Patient demonstrates a Factor XII mutation that is associated with the disease AND
2. Medications known to cause angioedema (i.e. ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued when appropriate AND
3. The requested agent will be used for prophylaxis against HAE attacks AND BOTH of the following:
 - a. Patient has a history of at least 2 acute attacks per month OR 2 severe attacks (laryngeal or abdominal attacks) per year OR for short-term prophylaxis prior to medical, surgical, or dental procedure AND
 - b. ONE of the following:
 - i. Patient has tried a formulary 17 alpha-alkylated androgen or antifibrinolytic agent OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a formulary 17 alpha-alkylated androgen or antifibrinolytic agent AND
4. ONE of the following:
 - a. Patient is receiving only ONE HAE agent indicated for prophylaxis against HAE attacks OR

- b. The other agent being used for prophylaxis will be discontinued before starting the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of hereditary angioedema (HAE) AND
3. The requested agent is being used for prophylaxis against HAE attacks AND
4. Patient has had a decrease in the frequency of acute attacks or has had stabilization of disease from use of the requested agent AND
5. ONE of the following:
 - a. Patient is receiving only ONE HAE agent indicated for prophylaxis against HAE attacks OR
 - b. The other agent being used for prophylaxis will be discontinued before starting the requested agent

Prior Authorization Group – Harvoni PA

Drug Name(s):

HARVONI

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The dosing of the requested agent is within the FDA labeled dosing or is AASLD/IDSA guideline supported

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group – Hetlioz PA

Drug Name(s):

HETLIOZ

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require the following:

1. Patient has a diagnosis of Non-24-Hour Sleep-Wake Disorder AND the patient is totally blind (i.e. no light perception)

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. neurologist, sleep specialist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – High Risk Medication PA - All Starts

Drug Name(s):

benztropine tablet
CLEMASTINE tablet
dicyclomine capsule, tablet
hydroxyzine hcl syrup, tablet
promethazine 12.5 mg, 25 mg suppository
promethazine syrup, tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

PA does NOT apply to patients less than 65 years of age.

Criteria for approval require ALL of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested high risk medication AND
2. Prescriber has indicated that the benefits of the requested high risk medication outweigh the risks for the patient AND
3. Prescriber has documented that s/he discussed risks and potential side effects of the requested high risk medication with the patient

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – High Risk Medication PA - Cyclobenzaprine

Drug Name(s):

cyclobenzaprine tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested medication AND
2. If patient is 65 years of age or over, then ALL of the following:
 - A. ONE of the following:
 - i. Patient has fibromyalgia and has tried and failed both duloxetine and Lyrica OR
 - ii. Patient has fibromyalgia and has a history of a documented intolerance, FDA labeled contraindication, or hypersensitivity to both duloxetine and Lyrica OR
 - iii. Patient has a diagnosis other than fibromyalgia which does NOT require any prerequisites AND
 - B. Prescriber has indicated that the benefits of the requested high risk medication outweigh the risks for the patient AND
 - C. Prescriber has documented that s/he discussed risks and potential side effects of the requested high risk medication with the patient

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – High Risk Medication PA - New Starts

Drug Name(s):

amitriptyline tablet

AMOXAPINE

clomipramine capsule

desipramine tablet

doxepin capsule, oral concentrate

imipramine hcl tablet

megestrol suspension, tablet

NORPRAMIN

nortriptyline capsule

NORTRIPTYLINE solution

paroxetine tablet

PAXIL

protriptyline tablet

SURMONTIL

trimipramine capsule

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only. PA does NOT apply to patients less than 65 years of age.

Criteria for approval require ONE of the following:

1. BOTH of the following:

A. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested high risk medication AND

B. There is evidence of a claim that the patient is currently being treated with the requested high risk medication within the past 180 days OR the prescriber states the patient is currently using the requested high risk medication OR

2. ALL of the following:

A. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested high risk medication AND

B. Prescriber has indicated that the benefits of the requested high risk medication outweigh the risks for the patient AND

C. Prescriber has documented that s/he discussed risks and potential side effects of the requested high risk medication with the patient

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – High Risk Medication PA - Zaleplon

Drug Name(s):

zaleplon capsule

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

PA does NOT apply to patients less than 65 years of age. Applies to greater than 90 days of therapy per calendar year.

Criteria for approval require ALL of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested high risk medication AND
2. Prescriber has indicated that the benefits of the requested high risk medication outweigh the risks for the patient AND
3. Prescriber has documented that s/he discussed risks and potential side effects of the requested high risk medication with the patient

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – High Risk Medication PA - Zolpidem

Drug Name(s):

zolpidem tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

PA does NOT apply to patients less than 65 years of age. Applies to greater than 90 days of therapy per calendar year.

Criteria for approval require ALL of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested high risk medication AND
2. Prescriber has indicated that the benefits of the requested high risk medication outweigh the risks for the patient AND
3. Prescriber has documented that s/he discussed risks and potential side effects of the requested high risk medication with the patient

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – HoFH PA - Juxtapid

Drug Name(s):

JUXTAPID

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of homozygous familial hypercholesterolemia (HoFH) AND
2. ONE of the following:
 - A. Genetic confirmation of two mutant alleles at the LDLR, Apo-B, PCSK9, ARH adaptor protein 1/LDLRAP1 gene locus OR
 - B. Untreated LDL-C greater than 500 mg/dL (greater than 13 mmol/L) or treated LDL-C 300 mg/dL or greater (7.76 mmol/L or greater) with ONE of the following:
 - i. Patient had cutaneous or tendon xanthoma before age 10 years OR
 - ii. Untreated elevated cholesterol levels consistent with heterozygous FH in both parents [untreated total cholesterol greater than 290 mg/dL (greater than 7.5 mmol/L) or untreated LDL-C greater than 190 mg/dL (greater than 4.9 mmol/L)] AND
3. ONE of the following:
 - A. Patient is on a maximally tolerated lipid-lowering regimen in the last 90 days (i.e. rosuvastatin in combination with ezetimibe OR atorvastatin in combination with ezetimibe) OR
 - B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a maximally tolerated lipid-lowering regimen (i.e. rosuvastatin in combination with ezetimibe OR atorvastatin in combination with ezetimibe) AND
4. Prescriber has taken baseline lab values for LDL-C, Apo-B, total cholesterol (TC), Non-HDL-C, and triglycerides (TG) AND
5. The requested agent will NOT be used in combination with Kynamro (mipomersen)

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of homozygous familial hypercholesterolemia (HoFH) AND
3. Patient has shown a reduction from baseline in at least ONE of the following metrics:
 - A. LDL-C OR
 - B. Apo-B OR
 - C. Total cholesterol (TC) OR
 - D. Non-HDL-C OR
 - E. Triglycerides (TG) AND
4. ONE of the following:
 - A. Patient is on a maximally tolerated lipid-lowering regimen in the last 90 days (i.e. rosuvastatin in combination with ezetimibe OR atorvastatin in combination with ezetimibe) OR
 - B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a maximally tolerated lipid-lowering regimen (i.e. rosuvastatin in combination with ezetimibe OR atorvastatin in combination with ezetimibe)
AND
5. The requested agent will NOT be used in combination with Kynamro (mipomersen)

Prior Authorization Group – HoFH PA - Kynamro

Drug Name(s):

KYNAMRO

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of homozygous familial hypercholesterolemia (HoFH) AND
2. ONE of the following:
 - A. Genetic confirmation of two mutant alleles at the LDLR, Apo-B, PCSK9, ARH adaptor protein 1/LDLRAP1 gene locus OR
 - B. Untreated LDL-C greater than 500 mg/dL (greater than 13 mmol/L) or treated LDL-C 300 mg/dL or greater (7.76 mmol/L or greater) with ONE of the following:
 - i. Patient had cutaneous or tendon xanthoma before age 10 years OR
 - ii. Untreated elevated cholesterol levels consistent with heterozygous FH in both parents [untreated total cholesterol greater than 290 mg/dL (greater than 7.5 mmol/L) or untreated LDL-C greater than 190 mg/dL (greater than 4.9 mmol/L)] AND
3. ONE of the following:
 - A. Patient is on a maximally tolerated lipid-lowering regimen in the last 90 days (i.e. rosuvastatin in combination with ezetimibe OR atorvastatin in combination with ezetimibe) OR
 - B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a maximally tolerated lipid-lowering regimen (i.e. rosuvastatin in combination with ezetimibe OR atorvastatin in combination with ezetimibe) AND
4. Prescriber has taken baseline lab values for LDL-C, Apo-B, total cholesterol (TC), Non-HDL-C, and triglycerides (TG) AND
5. The requested agent will NOT be used in combination with Juxtapid (Iomitapide)

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Initial approval 6 months. Renewal approval 12 months.

Other Criteria:

Criteria for renewal require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of homozygous familial hypercholesterolemia (HoFH) AND
3. Patient has shown a reduction from baseline in at least ONE of the following metrics:
 - A. LDL-C OR
 - B. Apo-B OR
 - C. Total cholesterol (TC) OR
 - D. Non-HDL-C OR
 - E. Triglycerides (TG) AND
4. ONE of the following:
 - A. Patient is on a maximally tolerated lipid-lowering regimen in the last 90 days (i.e. rosuvastatin in combination with ezetimibe OR atorvastatin in combination with ezetimibe) OR
 - B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a maximally tolerated lipid-lowering regimen (i.e. rosuvastatin in combination with ezetimibe OR atorvastatin in combination with ezetimibe)
AND
5. The requested agent will NOT be used in combination with Juxtapid (lomitapide)

Prior Authorization Group – Ilaris PA

Drug Name(s):

ILARIS

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D, plus the off-label use for acute gouty arthritis.

Exclusion Criteria:

An active or chronic infection (e.g. tuberculosis, HIV, hepatitis B/C)

Required Medical:

Criteria for approval require BOTH of:

1. ONE of:

- A. Patient (pt) has been diagnosed with Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS) OR
- B. Pt has been diagnosed with Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency (MKD) OR
- C. Pt has been diagnosed with Familial Mediterranean Fever (FMF) AND ONE of:
 - i. Pt has tried and failed colchicine OR
 - ii. Pt has a documented intolerance, FDA labeled contraindication, or hypersensitivity to colchicine OR
- D. Pt has been diagnosed with Cryopyrin-Associated Periodic Syndrome (CAPS) including Familial Cold Auto-inflammatory Syndrome (FCAS) or Muckle-Wells Syndrome (MWS) AND pt is at least 4 years of age OR
- E. Pt has been diagnosed with systemic juvenile idiopathic arthritis (SJIA) AND ALL of:
 - i. Pt is at least 2 years of age AND
 - ii. Pt has documented active systemic features (e.g. ongoing fever, anemia, rash, C-Reactive Protein levels greater than 50 mg/L, 2 or more joints with active arthritis) AND
 - iii. ONE of:
 - a. Pt has tried and failed at least ONE prerequisite agent (oral or IV glucocorticosteroids, prescription oral NSAIDs, methotrexate, leflunomide, Humira) OR
 - b. Pt has a documented intolerance, FDA labeled contraindication, or hypersensitivity to at least ONE prerequisite agent OR
- F. Pt has been diagnosed with acute gouty arthritis AND BOTH of:
 - i. Pt is at least 18 years of age AND
 - ii. ONE of:

- a. Pt has tried and failed at least TWO conventional first-line agents (prescription oral NSAIDs, colchicine, systemic corticosteroids) OR
 - b. Pt has a documented intolerance, FDA labeled contraindication, or hypersensitivity to at least TWO conventional first-line agents
- AND

2. ONE of:

- A. Pt is NOT currently being treated with another biologic agent OR
- B. Pt is currently being treated with another biologic agent and the agent will be discontinued prior to initiating the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Imiquimod PA

Drug Name(s):

ALDARA

imiquimod 5% cream

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently using the requested agent OR
 - c. Patient has ONE of the following diagnoses:
 - i. Actinic keratosis OR
 - ii. Superficial basal cell carcinoma
 - iii. External genital and/or perianal warts/condyloma acuminata OR
 - iv. Squamous cell carcinoma OR
 - v. Basal cell carcinoma OR
 - vi. Another indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

4 months for Actinic keratosis, other diagnoses - see Other Criteria

Other Criteria:

2 months for Superficial basal cell carcinoma, Squamous cell carcinoma, and Basal cell carcinoma

4 months for External genital and/or perianal warts/condyloma acuminata

12 months for All other diagnoses

Prior Authorization Group – Inbrija PA

Drug Name(s):

INBRIJA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require ALL of the following:

1. The requested agent will be used for intermittent treatment of OFF episodes in patients with Parkinson’s disease AND
2. Patient is receiving concurrent therapy with carbidopa/levodopa within the past 30 days AND
3. Patient is not currently taking a nonselective monoamine oxidase (MAO) inhibitor (e.g., phenelzine, tranylcypromine) or the patient has not recently (within 2 weeks) taken a nonselective MAO inhibitor

Age Restrictions:

Prescriber Restrictions:

Prescriber is a neurologist or the prescriber has consulted with a neurologist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Injectable Oncology PA

Drug Name(s):

ABRAXANE

ALIMTA

ALIQOPA

ARZERRA

AVASTIN

BELEODAQ

BESPONSA

BLINCYTO

CYRAMZA

DARZALEX

doxorubicin liposomal injection

EMPLICITI

ERBITUX

FOLOTYN

GAZYVA

HALAVEN

HERCEPTIN

HERCEPTIN HYLECTA

ISTODAX

JEVTANA

KADCYLA

KANJINTI

KYPROLIS

LARTRUVO

LUMOXITI

MVASI

MYLOTARG

ONIVYDE

PERJETA

POLIVY

PORTRAZZA

POTELIGEO

SYNRIBO

UNITUXIN

VECTIBIX

VELCADE

VYXEOS

YONDELIS

ZALTRAP

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.

Criteria for approval require BOTH of the following:

1. ONE of the following:

- A. Patient has an FDA labeled indication for the requested agent OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR
- C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. Patient has tried and failed the first line agent for the intended indication (if applicable) OR
 - b. Patient a documented intolerance, FDA labeled contraindication, or hypersensitivity to the first line agent AND
 - iii. Patient does NOT have any FDA labeled contraindication(s) to the requested agent AND
 - iv. Patient does NOT have any FDA labeled limitation(s) of use that is not otherwise supported in NCCN guidelines

May also be subject to Part B versus Part D review.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – IPF PA - Esbriet

Drug Name(s):

ESBRIET

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND
2. Patient has no known explanation for interstitial lung disease (ILD) or pulmonary fibrosis (e.g. radiation, drugs, metal dusts, sarcoidosis, or any connective tissue disease known to cause ILD) AND
3. ONE of the following:
 - A. Patient is NOT currently treated with Ofev (nintedanib) within the past 90 days
OR
 - B. Patient has been treated with Ofev (nintedanib) within the past 90 days AND therapy will be discontinued prior to starting therapy with the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND
3. The requested agent has been clinically beneficial to the patient AND
4. ONE of the following:
 - A. Patient is NOT currently treated with Ofev (nintedanib) within the past 90 days
OR
 - B. Patient has been treated with Ofev (nintedanib) within the past 90 days AND therapy will be discontinued prior to starting therapy with the requested agent

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. pulmonologist, radiologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – IPF PA - Ofev

Drug Name(s):

OFEV

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND
2. Patient has no known explanation for interstitial lung disease (ILD) or pulmonary fibrosis (e.g. radiation, drugs, metal dusts, sarcoidosis, or any connective tissue disease known to cause ILD) AND
3. ONE of the following:
 - A. Patient is NOT currently treated with Esbriet (pirfenidone) within the past 90 days OR
 - B. Patient has been treated with Esbriet (pirfenidone) within the past 90 days AND therapy will be discontinued prior to starting therapy with the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND
3. The requested agent has been clinically beneficial to the patient AND
4. ONE of the following:
 - A. Patient is NOT currently treated with Esbriet (pirfenidone) within the past 90 days OR
 - B. Patient has been treated with Esbriet (pirfenidone) within the past 90 days AND therapy will be discontinued prior to starting therapy with the requested agent

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. pulmonologist, radiologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Iron Chelating Agents PA - Exjade

Drug Name(s):

**deferasirox tablet
EXJADE**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ONE of the following:

- a. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome AND ONE of the following:
 - i. A liver iron (Fe) concentration (LIC) of at least 5 mg Fe per gram of dry weight OR
 - ii. A serum ferritin greater than 300 mcg/L OR
 - iii. MRI confirmation of iron deposition OR
- b. Patient has a diagnosis of chronic iron overload due to blood transfusions

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. ONE of the following:
 - a. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome OR
 - b. Patient has a diagnosis of chronic iron overload due to blood transfusions AND
3. Patient has shown clinical benefit with the requested agent

Age Restrictions:

For diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome, patient is at least 10 years of age. For diagnosis of chronic iron overload due to blood transfusions, patient is at least 2 years of age.

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Iron Chelating Agents PA - Jadenu

Drug Name(s):

**JADENU
JADENU SPRINKLE**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ONE of the following:

- a. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome AND ONE of the following:
 - i. A liver iron (Fe) concentration (LIC) of at least 5 mg Fe per gram of dry weight OR
 - ii. A serum ferritin greater than 300 mcg/L OR
 - iii. MRI confirmation of iron deposition OR
- b. Patient has a diagnosis of chronic iron overload due to blood transfusions

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. ONE of the following:
 - a. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome OR
 - b. Patient has a diagnosis of chronic iron overload due to blood transfusions AND
3. Patient has shown clinical benefit with the requested agent

Age Restrictions:

For diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome, patient is at least 10 years of age. For diagnosis of chronic iron overload due to blood transfusions, patient is at least 2 years of age.

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Kalydeco PA

Drug Name(s):

KALYDECO

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of cystic fibrosis AND
2. Patient has ONE of the CFTR gene mutations as indicated in the FDA label as confirmed by genetic testing AND
3. Patient is NOT homozygous for the F508del mutation AND
4. Patient has had pre-therapeutic/baseline FEV1 levels measured AND
5. ONE of the following:
 - A. Patient is NOT currently being treated with another CFTR agent [e.g. Orkambi (lumacaftor/ivacaftor), Symdeko (tezacaftor/ivacaftor and ivacaftor)] OR
 - B. Patient is currently being treated with another CFTR agent AND will discontinue the other CFTR agent prior to starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of cystic fibrosis AND
3. Patient has shown improvement or stabilization in FEV1 from pretreatment levels with the requested agent AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with another CFTR agent [e.g. Orkambi (lumacaftor/ivacaftor), Symdeko (tezacaftor/ivacaftor and ivacaftor)] OR
 - B. Patient is currently being treated with another CFTR agent AND will discontinue the other CFTR agent prior to starting the requested agent

Age Restrictions:

Patient is 6 months of age or over

Prescriber Restrictions:

Prescriber is a specialist (e.g. cystic fibrosis, pulmonologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Korlym PA

Drug Name(s):

KORLYM

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of Cushing's syndrome AND
2. ONE of the following:
 - A. Patient has type 2 diabetes mellitus OR
 - B. Patient has glucose intolerance (defined as 2-hour glucose tolerance test with glucose value of 140-199 mg/dL) AND
3. ONE of the following:
 - A. Patient has failed surgical resection OR
 - B. Patient is not a candidate for surgical resection

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Kuvan PA

Drug Name(s):

KUVAN

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of phenylketonuria (PKU) AND
2. Prescriber has submitted a baseline blood Phe level measured prior to initiation of therapy with the requested agent, which is above the recommended levels indicated for the patient's age range or condition AND
3. The dose is within FDA labeling

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of phenylketonuria (PKU) AND
3. ONE of the following:
 - a. Patient's blood Phe levels are being maintained within the acceptable range
 - OR
 - b. Patient has had a decrease in blood Phe level from baseline AND
4. The dose is within FDA labeling

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist with knowledge and expertise in metabolic diseases or genetic diseases or has consulted with a specialist in metabolic or genetic diseases

Coverage Duration:

Initial: 2 months if dose is 5 to less than 20 mg/kg/day, 1 month if 20 mg/kg/day

Renewal: 12 months

Other Criteria:

Prior Authorization Group – Lidocaine Topical PA - Lidocaine Gel/Jelly

Drug Name(s):

lidocaine 2% gel

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require the following:

1. The requested agent will be used for ONE of the following:
 - A. Surface anesthesia and lubrication for urethral procedure OR
 - B. Topical treatment for pain of urethritis OR
 - C. Surface anesthesia and lubrication for endotracheal intubation (oral and nasal)
OR
 - D. Another indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Lidocaine Topical PA - Lidocaine Ointment

Drug Name(s):

lidocaine 5% ointment

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require the following:

1. The requested agent will be used for ONE of the following:
 - A. Anesthesia of accessible mucous membranes of the oropharynx OR
 - B. Anesthetic lubricant for intubation OR
 - C. Temporary relief of pain associated with minor burns, including sunburn, abrasions of the skin, and insect bites OR
 - D. Another indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Lidocaine Topical PA - Lidocaine Patch

Drug Name(s):

**lidocaine 5% transdermal patch
LIDODERM**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require the following:

1. Patient has ONE of the following diagnoses:
 - A. Pain associated with postherpetic neuralgia (PHN) OR
 - B. Pain associated with diabetic neuropathy OR
 - C. Neuropathic pain associated with cancer OR
 - D. Another diagnosis that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Lidocaine Topical PA - Lidocaine Solution

Drug Name(s):

lidocaine 4% solution

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require the following:

1. The requested agent will be used for ONE of the following:
 - A. Topical anesthesia of accessible mucous membranes of the oral and nasal cavities OR
 - B. Topical anesthesia of accessible mucous membranes of proximal portions of the digestive tract OR
 - C. Another indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Lidocaine Topical PA - Lidocaine/prilocaine Cream

Drug Name(s):

lidocaine/prilocaine 2.5-2.5% cream

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require the following:

1. The requested agent will be used for ONE of the following:
 - A. Local analgesia on normal intact skin OR
 - B. Topical anesthetic for dermal procedures OR
 - C. Adjunctive anesthesia prior to local anesthetic infiltration in adult male genital skin OR
 - D. Anesthesia for minor procedures on female external genitalia OR
 - E. Another indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Linezolid PA

Drug Name(s):

linezolid suspension, tablet

ZYVOX suspension, tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. ONE of the following:

- a. The requested agent is prescribed by an infectious disease specialist or the prescriber has consulted with an infectious disease specialist on treatment of this patient AND the patient has an FDA labeled indication for the requested agent OR
- b. Patient has a documented infection due to vancomycin-resistant *Enterococcus faecium* OR
- c. Patient has a diagnosis of pneumonia caused by *Staphylococcus aureus* or *Streptococcus pneumoniae* AND ONE of the following:
 - i. Patient has a documented infection that is resistant to beta lactams, macrolides, clindamycin, tetracycline, and co-trimoxazole OR that is resistant to vancomycin OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to beta lactams, macrolides, clindamycin, tetracyclines, and co-trimoxazole OR
 - iii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to vancomycin OR
- d. Patient has a documented skin and skin structure infection, including diabetic foot infections, caused by *Staphylococcus aureus*, *Streptococcus pyogenes*, or *Streptococcus agalactiae* AND ONE of the following:
 - i. Patient has a documented infection that is resistant to beta lactams, macrolides, clindamycin, tetracycline, and co-trimoxazole OR that is resistant to vancomycin at the site of infection OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to beta lactams, macrolides, clindamycin, tetracyclines, and co-trimoxazole OR
 - iii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to vancomycin AND

2. ONE of the following:
 - a. Patient is NOT currently being treated for the same infection with Sivextro (tedizolid) OR
 - b. Current treatment with Sivextro (tedizolid) for the same infection will be discontinued before starting therapy with the requested agent AND
3. The dose is within the FDA labeled dosage

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 3 months

Other Criteria:

Prior Authorization Group – Linzess PA

Drug Name(s):

LINZESS

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. Patient has ONE of the following diagnoses:

A. Chronic idiopathic constipation with documentation of symptoms for at least 3 months OR

B. Irritable bowel syndrome with constipation with documentation of symptoms for at least 3 months AND

2. ONE of the following:

A. Patient has a medication history that includes lactulose or polyethylene glycol 3350 OR

B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to lactulose or polyethylene glycol 3350

Age Restrictions:

Patient is 18 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Mavyret PA

Drug Name(s):

MAVYRET

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The dosing of the requested agent is within the FDA labeled dosing or is AASLD/IDSA guideline supported AND
5. ONE of the following:
 - A. There is documentation that the patient is currently being treated with the requested agent OR
 - B. Patient has an FDA labeled contraindication or hypersensitivity to the preferred agents: Harvoni (ledipasvir/sofosbuvir) for supported genotypes OR Epclusa (sofosbuvir/velpatasvir) for genotype 2 or 3 OR
 - C. Prescriber has submitted documentation based on FDA approved labeling or AASLD/IDSA guidelines to support the use of the non-preferred agent for the patient's diagnosis and genotype over the preferred agents: Harvoni (ledipasvir/sofosbuvir) for supported genotypes OR Epclusa (sofosbuvir/velpatasvir) for genotype 2 or 3

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group – Memantine PA

Drug Name(s):

memantine solution, tablet
memantine titration pak
NAMENDA
NAMENDA TITRATION PAK

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

PA does NOT apply to patients 30 years of age or over

Criteria for approval require ONE of the following:

- A. Patient has a diagnosis of moderate to severe dementia of the Alzheimer's type OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Modafinil PA

Drug Name(s):

modafinil tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require BOTH of the following:

1. ONE of the following:
 - A. Patient has an FDA labeled indication for the requested agent OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - A. Patient is receiving only one of the listed agents, armodafinil OR modafinil, within the past 90 days OR
 - B. Patient has been treated with armodafinil within the past 90 days AND will discontinue prior to starting the requested agent

Age Restrictions:

Patient is 17 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – MS PA - Aubagio

Drug Name(s):

AUBAGIO

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. Patient is NOT currently being treated with a disease modifying agent (DMA) other than the requested agent OR
 - B. Patient is currently being treated with a DMA other than the requested agent AND the DMA will be discontinued before starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient is NOT currently being treated with an additional disease modifying agent (DMA) for the intended FDA labeled indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – MS PA - Avonex

Drug Name(s):

**AVONEX
AVONEX PREFILLED KIT**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. Patient is NOT currently being treated with a disease modifying agent (DMA) other than the requested agent OR
 - B. Patient is currently being treated with a DMA other than the requested agent AND the DMA will be discontinued before starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient is NOT currently being treated with an additional disease modifying agent (DMA) for the intended FDA labeled indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – MS PA - Betaseron

Drug Name(s):

BETASERON

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. Patient is NOT currently being treated with a disease modifying agent (DMA) other than the requested agent OR
 - B. Patient is currently being treated with a DMA other than the requested agent AND the DMA will be discontinued before starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient is NOT currently being treated with an additional disease modifying agent (DMA) for the intended FDA labeled indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – MS PA - Gilenya

Drug Name(s):

GILENYA 0.5 mg capsule

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. Patient is NOT currently being treated with a disease modifying agent (DMA) other than the requested agent OR
 - B. Patient is currently being treated with a DMA other than the requested agent AND the DMA will be discontinued before starting the requested agent AND
3. Prescriber has performed an electrocardiogram within 6 months prior to initiating treatment

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient is NOT currently being treated with an additional disease modifying agent (DMA) for the intended FDA labeled indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – MS PA - Glatiramer

Drug Name(s):

COPAXONE

glatiramer injection

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. Patient is NOT currently being treated with a disease modifying agent (DMA) other than the requested agent OR
 - B. Patient is currently being treated with a DMA other than the requested agent AND the DMA will be discontinued before starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient is NOT currently being treated with an additional disease modifying agent (DMA) for the intended FDA labeled indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – MS PA - Mayzent

Drug Name(s):

MAYZENT

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. Patient is NOT currently being treated with a disease modifying agent (DMA) other than the requested agent OR
 - B. Patient is currently being treated with a DMA other than the requested agent AND the DMA will be discontinued before starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient is NOT currently being treated with an additional disease modifying agent (DMA) for the intended FDA labeled indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – MS PA - Plegridy

Drug Name(s):

**PLEGRIDY
PLEGRIDY STARTER PACK**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. Patient is NOT currently being treated with a disease modifying agent (DMA) other than the requested agent OR
 - B. Patient is currently being treated with a DMA other than the requested agent AND the DMA will be discontinued before starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient is NOT currently being treated with an additional disease modifying agent (DMA) for the intended FDA labeled indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – MS PA - Tecfidera

Drug Name(s):

TECFIDERA

TECFIDERA STARTER PACK

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. Patient is NOT currently being treated with a disease modifying agent (DMA) other than the requested agent OR
 - B. Patient is currently being treated with a DMA other than the requested agent AND the DMA will be discontinued before starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient is NOT currently being treated with an additional disease modifying agent (DMA) for the intended FDA labeled indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – MS PA - Tysabri

Drug Name(s):

TYSABRI

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. Patient is NOT currently being treated with a disease modifying agent (DMA) other than the requested agent OR
 - B. Patient is currently being treated with a DMA other than the requested agent AND the DMA will be discontinued before starting the requested agent AND
3. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently using the requested agent AND is at risk if therapy is changed OR
 - C. Patient has ONE of the following diagnoses:
 - i. Multiple sclerosis (MS) AND ONE of the following:
 - a. Patient's medication history indicates the use of at least TWO preferred agents for the treatment of MS: Aubagio, Avonex, Betaseron, Gilenya, glatiramer (i.e. Copaxone, Glatopa), Mayzent, Plegridy, or Tecfidera OR
 - b. Patient has a documented intolerance, FDA labeled contraindication(s), or hypersensitivity to at least TWO preferred agents for the treatment of MS: Aubagio, Avonex, Betaseron, Gilenya, glatiramer (i.e. Copaxone, Glatopa), Mayzent, Plegridy, or Tecfidera OR
 - ii. Crohn's disease (CD) AND BOTH of the following:
 - a. ONE of the following:
 1. Patient's medication history indicates the use of at least ONE conventional CD therapy (e.g. aminosalicylates, metronidazole, ciprofloxacin, corticosteroids, methotrexate, or immunomodulators such as azathioprine or 6-mercaptopurine) OR

2. Patient has a documented intolerance, FDA labeled contraindication(s), or hypersensitivity to at least ONE conventional CD therapy AND
- b. ONE of the following:
1. Patient's medication history indicates use of ONE preferred biologic agent (Humira or Stelara) for the treatment of CD OR
 2. Patient has a documented intolerance, FDA labeled contraindication(s), or hypersensitivity to ONE preferred biologic agent (Humira or Stelara)

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months for MS, for CD 16 weeks for initial and 12 months renewal

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has shown clinical benefit with the requested agent AND
4. Patient is NOT currently being treated with an additional disease modifying agent (DMA) for the intended FDA labeled indication

Prior Authorization Group – Myalept PA

Drug Name(s):

MYALEPT

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of either congenital generalized lipodystrophy (CGL) or acquired generalized lipodystrophy (AGL) AND
2. Prescriber has drawn baseline values for HbA1C, triglycerides, and fasting insulin prior to beginning therapy with the requested agent AND
3. Patient also has at least ONE of the following additional diagnosis: diabetes mellitus, hypertriglyceridemia (200 mg/dL or higher), and/or high fasting insulin (30 μ U/mL or higher) AND
4. Patient has failed maximum tolerable dosing of a conventional agent for the additional diagnosis within the past 90 days AND
5. The dose is within the FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of either congenital generalized lipodystrophy (CGL) or acquired generalized lipodystrophy (AGL) AND
3. Patient has had a reduction in at least ONE of the following parameters: HbA1C, triglycerides and/or fasting insulin from baseline levels drawn prior to initiating the requested agent AND
4. The dose is within the FDA labeled dosing for the requested indication

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. endocrinologist, cardiologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Conventional agent examples include:

Hypertriglyceridemia: statins, fenofibrates, Omega-3-Acid Ethyl Esters (generic Lovaza)

Diabetes/high fasting insulin: insulin, sulfonylurea/sulfonylurea combination,
metformin/metformin combination

Prior Authorization Group – Natpara PA

Drug Name(s):

NATPARA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Increased baseline risk for osteosarcoma

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of hypocalcemia with hypoparathyroidism AND
2. Prescriber has confirmed the patient's 25-hydroxyvitamin D stores are not below the testing laboratory's normal range AND
3. Prescriber has confirmed the patient's serum calcium is above 7.5 mg/dL AND
4. ONE of the following:
 - A. Patient is NOT on concomitant use of alendronate OR
 - B. Patient is currently on alendronate and will discontinue it prior to therapy with the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of hypocalcemia with hypoparathyroidism AND
3. Patient has shown clinical benefit with the requested agent AND
4. Patient has an albumin-corrected total serum calcium concentration between 7.5 mg/dL and 10.6 mg/dL AND
5. ONE of the following:
 - A. Patient is NOT on concomitant use of alendronate OR
 - B. Patient is currently on alendronate and will discontinue it prior to therapy with the requested agent

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. endocrinologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Northera PA

Drug Name(s):

NORTHERA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of neurogenic orthostatic hypotension (NOH) AND
2. Prescriber has performed baseline blood pressure readings while the patient is sitting or supine (lying face up), AND also within 3 minutes of standing from a supine position AND
3. Patient has a decrease of at least 20 mmHg in systolic blood pressure or 10 mmHg diastolic blood pressure within 3 minutes after standing AND
4. Patient has persistent and consistent symptoms of neurogenic orthostatic hypotension (NOH) caused by ONE of the following:
 - A. Primary autonomic failure [Parkinson's disease (PD), multiple system atrophy, or pure autonomic failure] OR
 - B. Dopamine beta-hydroxylase deficiency OR
 - C. Non-diabetic autonomic neuropathy AND
5. Prescriber has assessed the severity of the patient's baseline symptoms of dizziness, lightheadedness, feeling faint, or feeling like the patient may black out

Criteria for renewal require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of neurogenic orthostatic hypotension (NOH) AND
3. Patient has demonstrated improvement in severity from baseline symptoms of dizziness, lightheadedness, feeling faint, or feeling like the patient may black out AND
4. Patient had an increase in systolic blood pressure from baseline of at least 10 mmHg upon standing from a supine (laying face up) position

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. cardiologist, neurologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be 1 month for initial, 3 months for renewal

Other Criteria:

Prior Authorization Group – Noxafil PA

Drug Name(s):

NOXAFIL

posaconazole 100 mg delayed release tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ONE of the following:

A. Patient has a diagnosis of oropharyngeal candidiasis AND patient has tried fluconazole or an alternative antifungal agent OR patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to fluconazole or an alternative antifungal agent OR

B. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or Candida AND patient is severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR

C. Patient has a diagnosis of invasive Aspergillus AND patient has tried an alternative antifungal agent OR patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to an alternative antifungal agent OR

D. Patient has another indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

One month for oropharyngeal candidiasis, 6 months for other indications

Other Criteria:

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. ONE of the following:

- A. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or Candida and patient continues to be severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR
- B. Patient has a diagnosis of invasive Aspergillus AND patient has continued indicators of active disease (e.g. continued radiologic findings, positive cultures, positive serum galactomannan assay for Aspergillus) OR
- C. Patient has another indication that is supported in CMS approved compendia for the requested agent

Prior Authorization Group – Nuedexta PA

Drug Name(s):

NUDEXTA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of pseudobulbar affect AND
2. Patient is NOT currently receiving a monoamine oxidase inhibitor (MAOI) OR the patient's MAOI will be discontinued at least 14 days prior to starting therapy with the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Nuplazid PA

Drug Name(s):

NUPLAZID

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Ocaliva PA

Drug Name(s):

OCALIVA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of Primary Biliary Cholangitis (PBC) as evidenced by TWO of the following three criteria at the time of diagnosis:

A. There is biochemical evidence of cholestasis with an alkaline phosphatase elevation of at least 1.5 times the upper limit of normal

B. Presence of antimitochondrial antibody (AMA): a titer greater than or equal to 1:40 OR a level that is above the testing laboratory's upper limit of the normal range

C. Histologic evidence of nonsuppurative destruction cholangitis and destruction of interlobular bile ducts AND

2. Prescriber has documented the patient's baseline (prior to any treatment with the requested agent) alkaline phosphatase (ALP) level AND total bilirubin level AND

3. ONE of the following:

A. BOTH of the following:

i. Patient has tried treatment with ursodiol and had an inadequate response AND

ii. Patient will continue treatment with ursodiol with the requested agent OR

B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ursodiol

Criteria for renewal require ALL of the following:

1. Patient has been previously approved for the requested agent through plan's PA criteria AND

2. Patient has a diagnosis of Primary Biliary Cholangitis (PBC) AND

3. ONE of the following:

A. Patient is currently on AND will continue treatment with ursodiol with the requested agent OR

B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ursodiol AND

4. Patient has had a decrease in alkaline phosphatase (ALP) level from baseline AND
5. Patient's total bilirubin is less than or equal to the upper limit of normal (ULN)

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Oncology Immunotherapy PA - Bavencio

Drug Name(s):

BAVENCIO

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.

Criteria for approval require BOTH of the following:

1. ONE of the following:

- A. Patient has an FDA labeled indication for the requested agent OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR
- C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. Patient has tried and failed the first line agent for the intended indication (if applicable) OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to the first line agent AND
 - iii. Patient does NOT have any FDA labeled limitation(s) of use that is not otherwise supported in NCCN guidelines

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Oncology Immunotherapy PA - Imfinzi

Drug Name(s):

IMFINZI

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.

Criteria for approval require BOTH of the following:

1. ONE of the following:

- A. Patient has an FDA labeled indication for the requested agent OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR
- C. ALL of the following:

- i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
- ii. ONE of the following:
 - a. Patient has tried and failed the first line agent for the intended indication (if applicable) OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to the first line agent AND
- iii. Patient does NOT have any FDA labeled limitation(s) of use that is not otherwise supported in NCCN guidelines

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Oncology Immunotherapy PA - Keytruda

Drug Name(s):

KEYTRUDA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.

Criteria for approval require BOTH of the following:

1. ONE of the following:

- A. Patient has an FDA labeled indication for the requested agent OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR
- C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. Patient has tried and failed the first line agent for the intended indication (if applicable) OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to the first line agent AND
 - iii. Patient does NOT have any FDA labeled limitation(s) of use that is not otherwise supported in NCCN guidelines

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Oncology Immunotherapy PA - Libtayo

Drug Name(s):

LIBTAYO

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.

Criteria for approval require BOTH of the following:

1. ONE of the following:

- A. Patient has an FDA labeled indication for the requested agent OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR
- C. ALL of the following:

- i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
- ii. ONE of the following:
 - a. Patient has tried and failed the first line agent for the intended indication (if applicable) OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to the first line agent AND
- iii. Patient does NOT have any FDA labeled limitation(s) of use that is not otherwise supported in NCCN guidelines

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Oncology Immunotherapy PA - Opdivo

Drug Name(s):

OPDIVO

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.

Criteria for approval require BOTH of the following:

1. ONE of the following:

- A. Patient has an FDA labeled indication for the requested agent OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR
- C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. Patient has tried and failed the first line agent for the intended indication (if applicable) OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to the first line agent AND
 - iii. Patient does NOT have any FDA labeled limitation(s) of use that is not otherwise supported in NCCN guidelines

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Oncology Immunotherapy PA - Tecentriq

Drug Name(s):

TECENTRIQ

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.

Criteria for approval require BOTH of the following:

1. ONE of the following:

- A. Patient has an FDA labeled indication for the requested agent OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR
- C. ALL of the following:

- i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
- ii. ONE of the following:
 - a. Patient has tried and failed the first line agent for the intended indication (if applicable) OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to the first line agent AND
- iii. Patient does NOT have any FDA labeled limitation(s) of use that is not otherwise supported in NCCN guidelines

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Oncology Immunotherapy PA - Yervoy

Drug Name(s):

YERVOY

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.

Criteria for approval require BOTH of the following:

1. ONE of the following:

- A. Patient has an FDA labeled indication for the requested agent OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR
- C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. Patient has tried and failed the first line agent for the intended indication (if applicable) OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to the first line agent AND
 - iii. Patient does NOT have any FDA labeled limitation(s) of use that is not otherwise supported in NCCN guidelines

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Ophthalmic Immunomodulators PA - Restasis

Drug Name(s):

RESTASIS

RESTASIS MULTIDOSE

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require ONE of the following:

1. Patient has an FDA labeled indication for the requested agent OR
2. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Opioids ER PA - Buprenorphine Pain

Drug Name(s):

**buprenorphine transdermal patch
BUTRANS**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - ii. Prescriber states the patient is currently using the requested agent within the past 180 days OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ALL of the following:
 - i. Prescriber provides documentation of a formal, consultative evaluation including:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - ii. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - iii. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - iv. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to immediate-acting opioid AND
 - v. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Opioids ER PA - Fentanyl Patch

Drug Name(s):

fentanyl transdermal patch

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - ii. Prescriber states the patient is currently using the requested agent within the past 180 days OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ALL of the following:
 - i. Prescriber provides documentation of a formal, consultative evaluation including:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - ii. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - iii. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - iv. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to immediate-acting opioid AND
 - v. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Opioids ER PA - Hydrocodone

Drug Name(s):

ZOHYDRO ER

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - ii. Prescriber states the patient is currently using the requested agent within the past 180 days OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ALL of the following:
 - i. Prescriber provides documentation of a formal, consultative evaluation including:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - ii. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - iii. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - iv. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to immediate-acting opioid AND
 - v. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Opioids ER PA - Morphine

Drug Name(s):

morphine sulfate ER tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - ii. Prescriber states the patient is currently using the requested agent within the past 180 days OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ALL of the following:
 - i. Prescriber provides documentation of a formal, consultative evaluation including:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - ii. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - iii. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - iv. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to immediate-acting opioid AND
 - v. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Opioids ER PA - Oxycodone

Drug Name(s):

OXYCONTIN

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - ii. Prescriber states the patient is currently using the requested agent within the past 180 days OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ALL of the following:
 - i. Prescriber provides documentation of a formal, consultative evaluation including:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - ii. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - iii. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - iv. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to immediate-acting opioid AND
 - v. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Opioids ER PA - Tapentadol

Drug Name(s):

NUCYNTA ER

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - ii. Prescriber states the patient is currently using the requested agent within the past 180 days OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ALL of the following:
 - i. Prescriber provides documentation of a formal, consultative evaluation including:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - ii. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - iii. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - iv. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to immediate-acting opioid AND
 - v. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Opioids ER PA - Tramadol

Drug Name(s):

tramadol ER tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for approval require ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - ii. Prescriber states the patient is currently using the requested agent within the past 180 days OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ALL of the following:
 - i. Prescriber provides documentation of a formal, consultative evaluation including:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - ii. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - iii. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - iv. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to immediate-acting opioid AND
 - v. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Oral Immunotherapy Agents PA - Grastek

Drug Name(s):

GRASTEK

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of allergic rhinitis, with or without conjunctivitis AND
2. Patient's diagnosis is confirmed with ONE of the following:
 - A. Positive skin test to ONE of the pollen extracts included in the requested agent: Timothy grass or cross-reactive grass OR
 - B. IgE specific antibodies to ONE of the extracts included in the requested agent: Timothy grass or cross-reactive grass AND
3. ONE of the following:
 - A. Patient has tried and failed at least TWO standard allergy medications, one of which was an intranasal corticosteroid OR
 - B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to at least TWO standard allergy medications, one of which was an intranasal corticosteroid AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with a subcutaneous injectable immunotherapy OR
 - B. Patient will discontinue subcutaneous injectable immunotherapy prior to starting the requested agent AND
5. ONE of the following:
 - A. Patient is NOT currently being treated with a beta blocker OR
 - B. Patient will discontinue the beta blocker prior to starting the requested agent AND
6. The requested agent will be started, or has already been started, 3 to 4 months before the expected onset of the applicable pollen season AND
7. The first dose is given in the clinic/hospital under direct supervision from the provider for a period of at least 30 minutes AND
8. Patient has been prescribed epinephrine auto-injector for at home emergency use

Age Restrictions:

Patient is between the ages of 5 and 65 years

Prescriber Restrictions:

Prescriber is a specialist (e.g. allergist, immunologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Standard allergy medications:

Oral antihistamines, oral corticosteroids, intranasal corticosteroids, intranasal antihistamines, or leukotriene inhibitors

Prior Authorization Group – Oral Immunotherapy Agents PA - Oralair

Drug Name(s):

ORALAIR

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of allergic rhinitis, with or without conjunctivitis AND
2. Patient's diagnosis is confirmed with ONE of the following:
 - A. Positive skin test to ONE of the pollen extracts included in the requested agent: Sweet vernal, orchard, perennial rye, Timothy, or Kentucky blue grass OR
 - B. IgE specific antibodies to ONE of the extracts included in the requested agent: Sweet vernal, orchard, perennial rye, Timothy, or Kentucky blue grass AND
3. ONE of the following:
 - A. Patient has tried and failed at least TWO standard allergy medications, one of which was an intranasal corticosteroid OR
 - B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to at least TWO standard allergy medications, one of which was an intranasal corticosteroid AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with a subcutaneous injectable immunotherapy OR
 - B. Patient will discontinue subcutaneous injectable immunotherapy prior to starting the requested agent AND
5. ONE of the following:
 - A. Patient is NOT currently being treated with a beta blocker OR
 - B. Patient will discontinue the beta blocker prior to starting the requested agent AND
6. The requested agent will be started, or has already been started, 3 to 4 months before the expected onset of the applicable pollen season AND
7. The first dose is given in the clinic/hospital under direct supervision from the provider for a period of at least 30 minutes AND
8. Patient has been prescribed epinephrine auto-injector for at home emergency use

Age Restrictions:

Patient is between the ages of 10 and 65 years

Prescriber Restrictions:

Prescriber is a specialist (e.g. allergist, immunologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Standard allergy medications:

Oral antihistamines, oral corticosteroids, intranasal corticosteroids, intranasal antihistamines, or leukotriene inhibitors

Prior Authorization Group – Oral Immunotherapy Agents PA - Ragwitek

Drug Name(s):

RAGWITEK

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of allergic rhinitis, with or without conjunctivitis AND
2. Patient's diagnosis is confirmed with ONE of the following:
 - A. Positive skin test to ONE of the pollen extracts included in the requested agent: Short Ragweed OR
 - B. IgE specific antibodies to ONE of the extracts included in the requested agent: Short Ragweed AND
3. ONE of the following:
 - A. Patient has tried and failed at least TWO standard allergy medications, one of which was an intranasal corticosteroid OR
 - B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to at least TWO standard allergy medications, one of which was an intranasal corticosteroid AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with a subcutaneous injectable immunotherapy OR
 - B. Patient will discontinue subcutaneous injectable immunotherapy prior to starting the requested agent AND
5. ONE of the following:
 - A. Patient is NOT currently being treated with a beta blocker OR
 - B. Patient will discontinue the beta blocker prior to starting the requested agent AND
6. The requested agent will be started, or has already been started, 3 to 4 months before the expected onset of the applicable pollen season AND
7. The first dose is given in the clinic/hospital under direct supervision from the provider for a period of at least 30 minutes AND
8. Patient has been prescribed epinephrine auto-injector for at home emergency use

Age Restrictions:

Patient is between the ages of 18 and 65 years

Prescriber Restrictions:

Prescriber is a specialist (e.g. allergist, immunologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Standard allergy medications:

Oral antihistamines, oral corticosteroids, intranasal corticosteroids, intranasal antihistamines, or leukotriene inhibitors

Prior Authorization Group – Orkambi PA

Drug Name(s):

ORKAMBI

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of cystic fibrosis AND
2. Patient has the presence of the F508del mutation on both alleles (homozygous) of the CFTR gene confirmed by genetic testing AND
3. Patient has had pre-therapeutic/baseline FEV1 levels measured AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with another CFTR agent [e.g. Kalydeco (ivacaftor), Symdeko (tezacaftor/ivacaftor and ivacaftor)] OR
 - B. Patient is currently being treated with another CFTR agent AND will discontinue the other CFTR agent prior to starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of cystic fibrosis AND
3. Patient has shown improvement or stabilization in FEV1 from pretreatment levels with the requested agent AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with another CFTR agent [e.g. Kalydeco (ivacaftor), Symdeko (tezacaftor/ivacaftor and ivacaftor)] OR
 - B. Patient is currently being treated with another CFTR agent AND will discontinue the other CFTR agent prior to starting the requested agent

Age Restrictions:

Patient is 2 years of age or over

Prescriber Restrictions:

Prescriber is a specialist (e.g. cystic fibrosis, pulmonologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Otezla PA

Drug Name(s):

OTEZLA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require BOTH of the following:

1. Patient has ONE of the following diagnoses:
 - A. Moderate-to-severe plaque psoriasis OR
 - B. Active psoriatic arthritis AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently using the requested agent AND is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of a biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's medication history indicates use of ONE conventional agent prerequisite for the requested indication OR
 - E. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to at least ONE conventional prerequisite agent for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of moderate-to-severe plaque psoriasis OR active psoriatic arthritis AND
3. Patient has shown clinical improvement (i.e. slowing of disease progression or decrease in symptom severity and/or frequency)

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Formulary conventional agent required for diagnoses of moderate-to-severe plaque psoriasis or active psoriatic arthritis

Formulary conventional agents for active psoriatic arthritis include methotrexate, hydroxychloroquine, sulfasalazine, minocycline, or leflunomide

Formulary conventional topical or systemic agents for plaque psoriasis include topical corticosteroids, tazarotene, cyclosporine, calcipotriene, methotrexate, tacrolimus, pimecrolimus, or acitretin

Prior Authorization Group – Palynziq PA

Drug Name(s):

PALYNZIQ

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of phenylketonuria (PKU) AND
2. Patient has a baseline blood Phe level greater than 600 micromol/L (10 mg/dL) AND
3. ONE of the following:
 - a. Patient is not also receiving Kuvan OR
 - b. Patient has been receiving Kuvan, AND will discontinue at least 14 days prior to receiving Palynziq AND
4. The dose is within the FDA labeled dose for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's prior authorization criteria AND
2. Patient has a diagnosis of phenylketonuria (PKU) AND
3. ONE of the following:
 - a. Patient's blood Phe levels are being maintained within the acceptable range OR
 - b. Patient has had a decrease in blood Phe level from baseline AND
4. ONE of the following:
 - a. Patient is not also receiving Kuvan OR
 - b. Patient has been receiving Kuvan, AND will discontinue at least 14 days prior to receiving Palynziq AND
5. The dose is within the FDA labeled dose for the requested indication

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist with knowledge and expertise in metabolic diseases or genetic diseases or has consulted with a specialist in metabolic or genetic diseases

Coverage Duration:

Initial approval 9 months. 12 months for renewal.

Other Criteria:

Prior Authorization Group – Pegylated Interferon PA

Drug Name(s):

PEGASYS

PEGASYS PROCLICK

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ONE of the following:

- A. Patient has a diagnosis of chronic hepatitis B AND BOTH of the following:
 - i. The chronic hepatitis B infection has been confirmed by serological markers AND
 - ii. Patient has NOT been administered the requested agent for more than 48 weeks for the treatment of chronic hepatitis B OR
- B. BOTH of the following:
 - i. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
 - ii. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype OR
- C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months for all other diagnoses. For hep B, hep C see Other Criteria

Other Criteria:

No prior peginterferon alfa use, approve 48 weeks for hepatitis B virus infection. Prior peginterferon alfa use, approve remainder of 48 weeks of total therapy for hepatitis B virus infection

Duration of therapy for hep C: Based on FDA approved labeling or AASLD/IDSA guideline supported

Prior Authorization Group – Prolia PA

Drug Name(s):

PROLIA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. ONE of the following:

A. Patient is a male or a postmenopausal female with a diagnosis of osteoporosis defined as ONE of the following:

i. Patient has a history of vertebral fracture(s), or low trauma or fragility fracture(s) [e.g. prior fracture from minor trauma such as falling from standing height or less] within the past 5 years OR

ii. Patient has a T-score that is -2.5 or lower AND ONE of the following:

1. Patient is female and medication history includes use of either a bisphosphonate or selective estrogen receptor (SERM) OR

2. Patient is male and medication history includes use of a bisphosphonate OR

3. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate or SERM (bisphosphonate or SERM for female patients, bisphosphonate for male patients) OR

B. Patient is requesting the agent for osteopenia (osteoporosis prophylaxis) AND ALL of the following:

i. ONE of the following:

1. Patient is a male 50 years of age and over OR

2. Patient is a postmenopausal female AND

ii. Patient has a T-score between -1.0 to -2.50 AND

iii. ONE of the following:

1. 10-year probability of a hip fracture 3% and greater per FRAX OR

2. 10-year probability of a major OP-related fracture 20% and greater per FRAX AND

iv. ONE of the following:

1. Patient is female and medication history includes use of a bisphosphonate or SERM OR

2. Patient is male and medication history includes use of a bisphosphonate OR
3. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate or SERM (bisphosphonate or SERM for female patients, bisphosphonate for male patients) OR

Criteria continues: See Other Criteria

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- C. Patient is a female with a diagnosis of breast cancer who is receiving aromatase inhibitor therapy AND ONE of:
 - i. Patient has a history of vertebral fracture(s), or low trauma or fragility fracture(s) [e.g. prior fracture from minor trauma such as falling from standing height or less] within the past 5 years OR
 - ii. Patient has a T-score at or below -1 AND ONE of:
 1. Patient's medication history includes use of a bisphosphonate OR
 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR
- D. Patient is a male with a diagnosis of prostate cancer receiving androgen deprivation therapy (ADT) AND ONE of:
 - i. Patient has a history of vertebral fracture(s), or low trauma or fragility fracture(s) [e.g. prior fracture from minor trauma such as falling from standing height or less] within the past 5 years OR
 - ii. BOTH of:
 1. ONE of:
 - a. Patient has a T-score at or below -1 OR
 - b. Patient has a history of an osteoporotic fracture AND
 2. ONE of:
 - a. Patient's medication history includes use of a bisphosphonate OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR
- E. Patient has a diagnosis of glucocorticoid-induced osteoporosis AND ALL of:
 - i. Patient is either initiating or continuing systemic glucocorticoids in a daily dose equivalent to 7.5 mg or greater of prednisone AND

- ii. Patient is expected to remain on glucocorticoids for at least 6 months AND
- iii. ONE of:
 - 1. Patient has a history of vertebral fracture(s), or low trauma or fragility fracture(s) [e.g. prior fracture from minor trauma such as falling from standing height or less] within the past 5 years OR
 - 2. Patient has a T-score that is -2.5 or lower AND ONE of:
 - a. Patient is female and medication history includes use of either a bisphosphonate or selective estrogen receptor (SERM) OR
 - b. Patient is male and medication history includes use of a bisphosphonate OR
 - c. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate or SERM (bisphosphonate or SERM for female patients, bisphosphonate for male patients) AND
- 2. ONE of:
 - A. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR
 - B. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR
 - C. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) AND
- 3. ONE of:
 - A. Patient is not receiving concomitant Forteo (teriparatide), Xgeva (denosumab), bisphosphonate, or SERM therapy within the past 90 days OR
 - B. Patient will discontinue the current Forteo (teriparatide), Xgeva (denosumab), bisphosphonate, or SERM therapy prior to initiating therapy with the requested agent AND
- 4. Dose requested is within the FDA approved labeling

Prior Authorization Group – Promacta PA

Drug Name(s):

PROMACTA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ONE of the following:

1. Patient (pt) has a diagnosis of chronic idiopathic thrombocytopenic purpura (ITP) AND ONE of the following:
 - A. Pt has had an insufficient response to single treatment with corticosteroids or immunoglobulins (IVIg or anti-D) OR
 - B. Pt has had an insufficient response to a splenectomy OR
 - C. BOTH of the following:
 - i. Pt is NOT a candidate for splenectomy AND
 - ii. Pt has a documented intolerance, FDA labeled contraindication or hypersensitivity to corticosteroids or immunoglobulins (IVIg or anti-D) OR
2. Pt has a diagnosis of hepatitis C associated thrombocytopenia AND ONE of the following:
 - A. Pt's platelet count is less than $75 \times 10^9/L$ AND the intent is to increase platelet counts sufficiently to initiate interferon therapy OR
 - B. Pt is on concurrent therapy with a pegylated interferon and ribavirin AND is at risk for discontinuing HCV therapy due to thrombocytopenia OR
3. Pt has a diagnosis of severe aplastic anemia (SAA) AND ALL of the following:
 - A. At least 2 of the following blood criteria:
 - i. Neutrophils less than $0.5 \times 10^9/L$ OR
 - ii. Platelets less than $20 \times 10^9/L$ OR
 - iii. Reticulocytes less than 1% corrected (percentage of actual hematocrit [Hct] to normal Hct) or reticulocyte count less than $20 \times 10^9/L$ AND
 - B. At least 1 of the following marrow criteria:
 - i. Severe hypocellularity is less than 25% OR
 - ii. Moderate hypocellularity is 25-50% with hematopoietic cells representing less than 30% of residual cells AND
 - C. ONE of the following:
 - i. Pt will be using the requested agent in combination with standard immunosuppressive therapy (i.e., antithymocyte globulin (ATG) and cyclosporine) for the first-line treatment of SAA (FSAA) OR
 - ii. ONE of following:

1. Pt has refractory (RSAA) defined as an insufficient response to immunosuppressive therapy [i.e., failure to ATG and cyclosporine] OR
2. Pt has a documented intolerance, FDA labeled contraindication, or hypersensitivity to horse ATG and cyclosporine

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Initial 6 months ITP & FSAA, 48 weeks HCV, 16 weeks RSAA Renewal 12 months ITP & SAA, 48 weeks HCV

Other Criteria:

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND

2. ONE of the following:

A. Patient has a diagnosis of chronic idiopathic thrombocytopenic purpura (ITP) AND ONE of the following:

- i. Patient's platelet count is $50 \times 10^9/L$ or greater OR
- ii. Patient's platelet count has increased sufficiently to avoid clinically important bleeding OR

B. Patient has a diagnosis of hepatitis C associated thrombocytopenia AND BOTH of the following:

i. ONE of the following:

1. Patient will be initiating hepatitis C therapy with interferon and ribavirin OR
2. Patient will be maintaining hepatitis C therapy with interferon and ribavirin at the same time as the requested agent AND

ii. ONE of the following:

1. Patient's platelet count is $90 \times 10^9/L$ or greater OR
2. Patient's platelet count has increased sufficiently to initiate or maintain interferon based therapy for the treatment of hepatitis C OR

C. Patient has a diagnosis of refractory severe aplastic anemia (SAA) AND has had a hematological response by week 16 defined as ONE of the following:

- i. Platelet count increases to $20 \times 10^9/L$ above baseline OR
- ii. Stable platelet counts with transfusion independence for a minimum of 8 weeks OR
- iii. Hemoglobin increase by greater than 1.5 g/dL OR

- iv. Reduction in 4 units or greater of red blood cell (RBC) transfusions for 8 consecutive weeks OR
 - v. An absolute neutrophil count (ANC) increase of 100% OR
 - vi. An absolute neutrophil count (ANC) increase greater than $0.5 \times 10^9/L$ OR
- D. Patient has a diagnosis of severe aplastic anemia AND BOTH of the following:
- i. Patient is using the requested agent in combination with standard immunosuppressive therapy (i.e., antithymocyte globulin (ATG) and cyclosporine) for the first-line treatment of severe aplastic anemia AND
 - ii. ONE of the following:
 - 1. Patient has had a complete response by 6 months defined as hematological parameters meeting ALL of the following values:
 - a) An absolute neutrophil count (ANC) greater than 1,000/mcL AND
 - b) Platelet count greater than $100 \times 10^9/L$ AND
 - c) Hemoglobin greater than 10 g/dL OR
 - 2. Patient has had a partial response by 6 months defined as meeting TWO of the following values:
 - a) An absolute neutrophil count (ANC) greater than 500/mcL OR
 - b) Platelet count greater than $20 \times 10^9/L$ OR
 - c) Reticulocyte count greater than 60,000/mcL

Prior Authorization Group – Pulmonary Hypertension PA - Adcirca

Drug Name(s):

ADCIRCA

tadalafil 20 mg tablet (generic for Adcirca)

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent AND Concurrently taking an phosphodiesterase type 5 (PDE-5) inhibitor (e.g. Cialis, Viagra) with the requested agent

Required Medical:

Criteria for initial approval require ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient (pt) is currently being treated with the requested agent within the past 90 days OR
- b. Prescriber states pt is using the requested agent AND is at risk if therapy is changed AND

ii. Pt has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. Pt has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Pt's World Health Organization (WHO) functional class is II or greater AND

ii. Pt has a mean pulmonary arterial pressure greater than or equal to 25 mmHg AND

iii. Pt has a pulmonary vascular resistance greater than 3 Wood units AND

iv. ONE of the following:

- a. Requested agent will be utilized as monotherapy OR
- b. Requested agent is for use in combination with ambrisentan for dual therapy ONLY OR
- c. Requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy: except for dual therapy requests for tadalafil with ambrisentan), then BOTH of the following:

1. Pt has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

2. Requested agent is in a different therapeutic class OR

d. Requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:

1. A prostanoid has been started as one of the agents in the triple therapy unless the pt has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a prostanoid AND
2. Pt has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
3. All three agents in the triple therapy are from a different therapeutic class OR

C. Pt has an indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Pt has been previously approved for the requested agent through the plan's PA criteria AND
2. Pt has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
3. Pt is responding to therapy with the requested agent

Prior Authorization Group – Pulmonary Hypertension PA - Adempas

Drug Name(s):

ADEMPAS

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require BOTH of the following:

1. ONE of:

A. BOTH of:

i. ONE of:

- a. There is evidence of a claim that patient (pt) is currently being treated with the requested agent within the past 90 days OR
- b. Prescriber states the pt is using the requested agent AND is at risk if therapy is changed AND

ii. Pt has an FDA labeled indication for the requested agent OR

B. Pt has a diagnosis (dx) of CTEPH, WHO Group 4 as determined by ventilation-perfusion scan and a confirmatory selective pulmonary angiography AND ALL of:

i. ONE of:

- a. Pt is NOT a candidate for surgery OR
- b. Pt has had pulmonary endarterectomy AND has persistent or recurrent disease AND

ii. Pt has a mean pulmonary arterial pressure greater than or equal to 25 mmHg AND

iii. Pt has a pulmonary capillary wedge pressure less than or equal to 15 mmHg OR

C. Pt has a dx of PAH, WHO Group 1 as determined by right heart catheterization AND ALL of:

i. Pt's WHO functional class is II or greater AND

ii. Pt has a mean pulmonary arterial pressure greater than or equal to 25 mmHg AND

iii. Pt has a pulmonary vascular resistance greater than 3 Wood units AND

2. ONE of:

A. Requested agent will be utilized as monotherapy OR

B. Requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), then BOTH of:

- i. Pt has unacceptable or deteriorating clinical status despite established pharmacotherapy AND
 - ii. Requested agent is in a different therapeutic class OR
- C. Requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), then ALL of:
 - i. Prostanoid therapy has been started as one of the agents in the triple therapy unless the pt has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a prostanoid AND
 - ii. Pt has unacceptable or deteriorating clinical status despite established pharmacotherapy AND
 - iii. All three agents in the triple therapy are from a different therapeutic class

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Pt has been previously approved for the requested agent through the plan's PA criteria AND
2. Pt has an FDA labeled indication for the requested agent AND
3. Pt is responding to therapy with the requested agent

Prior Authorization Group – Pulmonary Hypertension PA - Bosentan

Drug Name(s):

bosentan tablet
TRACLEER

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Elevated liver enzymes accompanied by signs or symptoms of liver dysfunction/injury or a bilirubin level of 2 times the ULN (upper limit of normal) or greater AND FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ONE of the following:

1. BOTH of the following:
 - A. ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is using the requested agent AND is at risk if therapy is changed AND
 - B. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR
2. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1, as determined by right heart catheterization, AND ALL of the following:
 - i. Patient's WHO functional class is II or greater AND
 - ii. Patient has a mean pulmonary arterial pressure greater than or equal to 25 mmHg AND
 - iii. Patient has a pulmonary vascular resistance greater than 3 Wood units AND
 - iv. ONE of the following:
 - a. Requested agent will be utilized as monotherapy OR
 - b. Requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), then BOTH of the following:
 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 2. Requested agent is in a different therapeutic class OR
 - c. Requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), then ALL of the following:

1. A prostanoid has been started as one of the agents in the triple therapy unless the patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a prostanoid AND
 2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 3. All three agents in the triple therapy are from a different therapeutic class OR
3. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
3. Patient is responding to therapy with the requested agent

Prior Authorization Group – Pulmonary Hypertension PA - Letairis

Drug Name(s):

ambrisentan tablet

LETAIRIS

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
- b. Prescriber states the patient is using the requested agent AND is at risk if therapy is changed AND

ii. Patient has an FDA labeled indication for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Patient's WHO functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than or equal to 25 mmHg AND

iii. Patient has a pulmonary vascular resistance greater than 3 Wood units AND

iv. ONE of the following:

- a. Requested agent will be utilized as monotherapy OR
- b. Requested agent is for use in combination with tadalafil for dual therapy ONLY OR

c. Requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy: except dual therapy requests for ambrisentan with tadalafil), then BOTH of the following:

- 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
- 2. Requested agent is in a different therapeutic class OR

d. Requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), then ALL of the following:

- 1. A prostanoid has been started as one of the agents in the triple therapy unless the patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
3. All three agents in the triple therapy are from a different therapeutic class

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient is responding to therapy with the requested agent

Prior Authorization Group – Pulmonary Hypertension PA - Opsumit

Drug Name(s):

OPSUMIT

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
- b. Prescriber states the patient is using the requested agent AND is at risk if therapy is changed AND

ii. Patient has an FDA labeled indication for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Patient's WHO functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than or equal to 25 mmHg AND

iii. Patient has a pulmonary vascular resistance greater than 3 Wood units AND

iv. ONE of the following:

a. Requested agent will be utilized as monotherapy OR

b. Requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), then BOTH of the following:

1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

2. Requested agent is in a different therapeutic class OR

c. Requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), then ALL of the following:

1. A prostanoid has been started as one of the agents in the triple therapy unless the patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

3. All three agents in the triple therapy are from a different therapeutic class

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient is responding to therapy with the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Pulmonary Hypertension PA - Sildenafil

Drug Name(s):

sildenafil 20 mg tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent AND Concurrently taking an phosphodiesterase type 5 (PDE-5) inhibitor (e.g. Cialis, Viagra) with the requested agent

Required Medical:

Criteria for initial approval require ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
- b. Prescriber states the patient is using the requested agent AND is at risk if therapy is changed AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

- i. Patient's World Health Organization (WHO) functional class is II or greater AND
- ii. Patient has a mean pulmonary arterial pressure greater than or equal to 25 mmHg AND
- iii. Patient has a pulmonary vascular resistance greater than 3 Wood units AND
- iv. ONE of the following:

- a. Requested agent will be utilized as monotherapy OR
- b. Requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), then BOTH of the following:
 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 2. Requested agent is in a different therapeutic class OR
- c. Requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:
 1. A prostanoid has been started as one of the agents in the triple therapy unless the patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
3. All three agents in the triple therapy are from a different therapeutic class OR

C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
3. Patient is responding to therapy with the requested agent

Prior Authorization Group – Pulmonary Hypertension PA - Uptravi

Drug Name(s):

UPTRAVI

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
- b. Prescriber states the patient is using the requested agent AND is at risk if therapy is changed AND

ii. Patient has an FDA labeled indication for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following

- i. Patient's World Health Organization (WHO) functional class is II or greater AND
- ii. Patient has a mean pulmonary arterial pressure greater than or equal to 25 mmHg AND

iii. Patient has a pulmonary vascular resistance greater than 3 Wood units AND

iv. ONE of the following:

- a. Requested agent will be utilized as monotherapy OR
- b. Requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), then BOTH of the following:
 - 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 2. Requested agent is in a different therapeutic class OR
- c. Requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), then BOTH of the following:
 - 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 2. All three agents in the triple therapy are from a different therapeutic class

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient is responding to therapy with the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Pulmonary Hypertension PA - Ventavis

Drug Name(s):

VENTAVIS

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
- b. Prescriber states the patient is using the requested agent AND is at risk if therapy is changed AND

ii. Patient has an FDA labeled indication for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following

- i. Patient's World Health Organization (WHO) functional class is II or greater AND
- ii. Patient has a mean pulmonary arterial pressure greater than or equal to 25 mmHg AND

iii. Patient has a pulmonary vascular resistance greater than 3 Wood units AND

iv. ONE of the following:

- a. Requested agent will be utilized as monotherapy OR
- b. Requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), then BOTH of the following:
 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 2. Requested agent is in a different therapeutic class OR
- c. Requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), then ALL of the following:
 1. Patient is WHO functional class III or IV AND
 2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 3. All three agents in the triple therapy are from a different therapeutic class

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient is responding to therapy with the requested agent

Drug is also subject to Part B versus Part D review.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Regranex PA

Drug Name(s):

REGRANEX

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ONE of the following:

1. BOTH of the following:

A. Patient has a diagnosis of lower extremity diabetic neuropathic ulcer(s) that extend into the subcutaneous tissue or beyond AND

B. The ulcer(s) intended for treatment has an adequate blood supply OR

2. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Relistor PA

Drug Name(s):

RELISTOR

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. ONE of the following diagnoses:
 - A. Patient has opioid-induced constipation (OIC) with advanced illness or pain caused by active cancer and is receiving palliative care AND the requested agent is Relistor (methylnaltrexone) injection OR
 - B. Patient has opioid-induced constipation (OIC) and chronic non-cancer pain, including patients with chronic pain related to prior cancer or its treatment AND
2. Patient has chronic use of an opioid agent in the past 90 days AND
3. ONE of the following:
 - A. Patient has a medication history that includes lactulose or polyethylene glycol 3350 OR
 - B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to lactulose or polyethylene glycol 3350

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Repatha PA

Drug Name(s):

REPATHA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA approved diagnosis for the requested agent AND
2. ONE of the following:
 - A. If the diagnosis is heterozygous familial hypercholesterolemia (HeFH), ONE of the following:
 - i. Genetic confirmation of one mutant allele at the LDLR, Apo-B, PCSK9, or ARH adaptor protein 1/LDLRAP1 gene locus OR
 - ii. BOTH of the following:
 1. Total cholesterol greater than 290 mg/dL (greater than 7.5 mmol/L) (pretreatment or highest level while on treatment) OR LDL-C greater than 190 mg/dL (greater than 4.9 mmol/L) (pretreatment or highest level while on treatment) AND
 2. History of tendon xanthomas in ONE of the following:
 - a. Patient
 - b. Patient's first degree relative (i.e. parent, sibling, or child)
 - c. Patient's second degree relative (e.g. grandparent, uncle, or aunt) OR
 - iii. Patient has a Dutch Lipid Clinic Network Criteria score of greater than 8 OR
 - B. If the diagnosis is homozygous familial hypercholesterolemia (HoFH), ONE of the following:
 - i. Genetic confirmation of two mutant alleles at the LDLR, Apo-B, PCSK9, or ARH adaptor protein 1/LDLRAP1 gene locus OR
 - ii. Untreated LDL-C greater than 500 mg/dL (greater than 13 mmol/L) or treated LDL-C 300 mg/dL or greater (7.76 mmol/L or greater) with ONE of the following:
 - a. Patient had cutaneous or tendon xanthoma before age 10 years OR
 - b. Untreated elevated cholesterol levels consistent with heterozygous FH in both parents [untreated LDL-C greater than

190 mg/dL (greater than 4.9 mmol/L) or untreated total cholesterol greater than 290 mg/dL (greater than 7.5 mmol/L)] OR

Initial criteria continues: see Other Criteria

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. cardiologist or endocrinologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- C. Patient's indication is to reduce the risk of myocardial infarction, stroke, and coronary revascularization in patients with established cardiovascular disease (angina pectoris, coronary heart disease, myocardial infarction, transient ischemic attacks, cerebrovascular disease (CeVD) or peripheral vascular disease (PVD) or after coronary revascularization or carotid endarterectomy) AND
3. ONE of the following:
- A. ONE of the following:
 - 1. Patient is currently (for the past 90 days) on high-intensity statin therapy (i.e. rosuvastatin 20-40 mg or atorvastatin 40-80 mg) OR
 - 2. BOTH of the following:
 - a. Patient has tried and is intolerant to high-intensity statin therapy AND
 - b. Patient is currently (for the past 90 days) on low or moderate intensity statin therapy OR
 - B. Patient has documented intolerance* to TWO different statins (*intolerance is defined as inability to tolerate the lowest FDA approved starting dose of a statin) OR
 - C. Patient has an FDA labeled contraindication to a statin AND
4. Praluent (alirocumab) and Repatha (evolocumab) will not be used concurrently with each other

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
- 2. Patient has an FDA approved diagnosis for the requested agent AND

3. Patient has shown clinical benefit with the requested agent AND
4. ONE of the following:
 - A. ONE of the following:
 1. Patient is currently (for the past 90 days) on high-intensity statin therapy (i.e. rosuvastatin 20-40 mg or atorvastatin 40-80 mg) OR
 2. BOTH of the following:
 - a. Patient has tried and is intolerant to high-intensity statin therapy AND
 - b. Patient is currently (for the past 90 days) on low or moderate intensity statin therapy OR
 - B. Patient has documented intolerance* to TWO different statins (*intolerance is defined as inability to tolerate the lowest FDA approved starting dose of a statin) OR
 - C. Patient has an FDA labeled contraindication to a statin AND
5. Praluent (alirocumab) and Repatha (evolocumab) will not be used concurrently with each other

Prior Authorization Group – Samsca PA

Drug Name(s):

SAMSCA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the request agent AND Any underlying liver disease, including cirrhosis

Required Medical:

Criteria for approval require ALL of the following:

1. The requested agent was initiated (or re-initiated) in the hospital AND
2. Prior to initiating the requested agent, patient has or had a diagnosis of clinically significant hypervolemic or euvolemic hyponatremia defined by ONE of the following:
 - a. Serum sodium is less than 125 mEq/L OR
 - b. Serum sodium is 125 mEq/L or greater AND patient has symptomatic hyponatremia that has resisted correction with fluid restriction AND
3. Medications known to cause hyponatremia have been evaluated and discontinued when appropriate AND
4. Patient has NOT already received 30 days of therapy with the requested agent following the most recent hospitalization for initiation of therapy AND
5. Requested dose is within the FDA approved labeled dosing (Initial dose is 15 mg once daily, may be increased to 30 mg once daily after 24 hours, up to a maximum daily dose of 60 mg, as needed to achieve the desired level of serum sodium. Do not administer the requested agent for more than 30 days to minimize the risk of liver injury)

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 30 days

Other Criteria:

Prior Authorization Group – Self - Administered Oncology PA

Drug Name(s):

abiraterone 250 mg tablet

AFINITOR

AFINITOR DISPERZ

ALECENSA

ALUNBRIG

BALVERSA

bexarotene capsule

BOSULIF

BRAFTOVI

CABOMETYX

CALQUENCE

CAPRELSA

COMETRIQ

COPIKTRA

COTELLIC

DAURISMO

ERIVEDGE

ERLEADA

erlotinib tablet

FARYDAK

GILOTRIF

GLEEVEC

HEXALEN

IBRANCE

ICLUSIG

IDHIFA

imatinib mesylate tablet

IMBRUVICA

INLYTA

IRESSA

JAKAFI

KISQALI

KISQALI FEMARA CO PACK

LENVIMA

LONSURF

LORBRENA

LYNPARZA

MATULANE

MEKINIST

MEKTOVI

NERLYNX

NEXAVAR

NINLARO

ODOMZO

PIQRAY
POMALYST
REVLIMID
RUBRACA
RYDAPT
SPRYCEL
STIVARGA
SUTENT
SYLATRON
TAFINLAR
TAGRISSO
TALZENNA
TARCEVA
TARGRETIN 75 mg capsule
TASIGNA
THALOMID
TIBSOVO
tretinoin capsule
TYKERB
VENCLEXTA
VENCLEXTA STARTING PACK
VERZENIO
VITRAKVI
VIZIMPRO
VOTRIENT
XALKORI
XOSPATA
XPOVIO
XTANDI
YONSA
ZEJULA
ZELBORAF
ZOLINZA
ZYDELIG
ZYKADIA
ZYTIGA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.
Criteria for approval require BOTH of the following:

1. ONE of the following:
 - A. Patient has an FDA labeled indication for the requested agent OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. Patient does NOT have any FDA labeled contraindication(s) to the requested agent AND
 - iii. ONE of the following:
 1. Patient has tried and failed the first line agent for the intended indication (if applicable) OR
 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to the first line agent AND
 - iv. Patient does NOT have any FDA labeled limitation(s) of use that is not otherwise supported in NCCN guidelines

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Signifor LAR PA

Drug Name(s):

SIGNIFOR LAR

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Severe hepatic impairment (i.e. Child Pugh C)

Required Medical:

Criteria for initial approval require ONE of the following:

1. Patient has a diagnosis of acromegaly AND ONE of the following:
 - i. Patient had an inadequate response to surgery defined by ONE of the following:
 - a. Growth hormone level greater than 5 ng/mL OR
 - b. IGF-1 level greater than 1.9 U/mL for males or greater than 2.2 U/mL for females OR
 - ii. Patient is NOT a candidate for surgical resection OR
2. Patient has a diagnosis of Cushing's disease AND ONE of the following:
 - i. Patient has recurrence or persistence of symptoms after pituitary surgical resection OR
 - ii. Patient is NOT a candidate for pituitary surgical resection OR
3. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. ONE of the following:
 - A. Patient has a diagnosis of acromegaly AND ONE of the following:
 - i. Growth hormone level less than 5 ng/mL OR
 - ii. IGF-1 level less than 1.9 U/mL for males or less than 2.2 U/mL for females OR
 - iii. Clinical improvement (e.g. reduction in tumor size, decreased headaches, improved cardiovascular or respiratory symptoms) OR
 - B. Patient has a diagnosis of Cushing's disease AND BOTH of the following:
 - i. Patient has a urinary free cortisol level less than or equal to the upper limit of normal AND
 - ii. Patient has shown improvement in at least ONE of the following clinical signs and symptoms
 1. Fasting plasma glucose OR
 2. Hemoglobin A1c OR
 3. Hypertension OR

4. Weight OR

C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Initial: Acromegaly - 6 months, Cushing - 7 months, All other dxs - 12 months, Renewal: 12 months

Other Criteria:

Prior Authorization Group – Signifor PA

Drug Name(s):

SIGNIFOR

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Severe hepatic impairment (i.e. Child Pugh C)

Required Medical:

Criteria for initial approval require ONE of the following:

1. Patient has a diagnosis of Cushing's disease (CD) AND ONE of the following:
 - i. Patient has had recurrence or persistence of symptoms after pituitary surgical resection OR
 - ii. Patient is NOT a candidate for pituitary surgical resection OR
2. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. ONE of the following:
 - A. Patient has a diagnosis of Cushing's disease (CD) AND BOTH of the following:
 - i. Patient has had a 15% or greater decrease in urinary free cortisol levels AND
 - ii. Patient has shown improvement in at least ONE of the following clinical signs and symptoms:
 1. Fasting plasma glucose OR
 2. Hemoglobin A1c OR
 3. Hypertension OR
 4. Weight OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Initial approval: 3 months for CD, 12 months for all other diagnoses, Renewal approval: 12 months

Other Criteria:

Prior Authorization Group – Sivextro PA

Drug Name(s):

SIVEXTRO tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following:
 - a. BOTH of the following:
 - i. A documented acute bacterial skin and skin structure infection (ABSSSI) defined as a bacterial infection of the skin with a lesion size area of at least 75 cm² (lesion size measured by the area of redness, edema, or induration) AND
 - ii. The infection is due to *Staphylococcus aureus*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus anginosus*, *Streptococcus intermedius*, *Streptococcus constellatus*, or *Enterococcus faecalis* OR
 - b. Another indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - a. The requested agent is prescribed by an infectious disease specialist or the prescriber has consulted with an infectious disease specialist on treatment of this patient OR
 - b. The requested agent is NOT prescribed by an infectious disease specialist or the prescriber has NOT consulted with an infectious disease specialist on treatment of this patient AND ONE of the following:
 - i. There is documentation of resistance to beta lactams, macrolides, clindamycin, tetracycline, and co-trimoxazole at the site of infection OR the patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to beta lactams, macrolides, clindamycin, tetracyclines, and co-trimoxazole OR
 - ii. There is documentation of resistance to vancomycin at the site of infection OR the patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to vancomycin AND
3. ONE of the following:
 - a. Patient is NOT currently being treated for the same infection with linezolid OR
 - b. The current treatment with linezolid for the same infection will be discontinued before starting therapy with the requested agent AND
4. The requested dose is within the FDA and/or compendia labeled dosage

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be 6 days for ABSSSI or 30 days for all other indications

Other Criteria:

Prior Authorization Group – Somatostatin Analogs PA - Octreotide

Drug Name(s):

octreotide injection

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require BOTH of the following:

1. ONE of the following:

- A. Patient has an FDA approved indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently using the requested agent AND is at risk if therapy is changed OR

B. ONE of the following:

- i. Patient has a diagnosis of acromegaly AND ONE of the following:
 - a. Patient is not a candidate for surgical resection or pituitary radiation therapy OR
 - b. The requested agent is for adjunctive therapy with pituitary radiation therapy OR
 - c. Patient had an inadequate response to surgery or pituitary radiation therapy as indicated by growth hormone levels or serum IGF-1 levels that are above the reference range OR
- ii. Patient has severe diarrhea and/or flushing episodes associated with metastatic carcinoid tumors OR
- iii. Patient has profuse watery diarrhea associated with Vasoactive Intestinal Peptide (VIP) secreting tumors OR
- iv. Patient has a diagnosis of dumping syndrome AND the following:
 - a. Patient has tried acarbose or has a documented intolerance, FDA labeled contraindication, or hypersensitivity to acarbose OR
 - v. Patient has another indication that is supported in CMS approved compendia for the requested agent AND

2. The dose requested is within the FDA approved or CMS approved compendia dosing for the requested agent and indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Initial approval will be for 6 months, renewal approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria and has responded to therapy AND
2. ONE of the following:
 - A. Patient has a diagnosis of acromegaly AND the following:
 - i. Patient has had clinical improvement (e.g. reduction in tumor size, decreased headaches, improved cardiovascular or respiratory symptoms)
OR
 - B. Patient has severe diarrhea and/or flushing episodes associated with metastatic carcinoid tumors OR
 - C. Patient has profuse watery diarrhea associated with Vasoactive Intestinal Peptide (VIP) secreting tumors OR
 - D. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
3. The dose requested is within the FDA approved or CMS approved compendia dosing for the requested agent and indication

Prior Authorization Group – Somatostatin Analogs PA - Sandostatin LAR

Drug Name(s):

SANDOSTATIN LAR

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. ONE of the following:

- A. Patient has an FDA approved indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently using the requested agent AND is at risk if therapy is changed OR

B. ONE of the following:

- i. Patient has a diagnosis of acromegaly AND ONE of the following:
 - a. Patient is not a candidate for surgical resection or pituitary radiation therapy OR
 - b. The requested agent is for adjunctive therapy with pituitary radiation therapy OR
 - c. Patient had an inadequate response to surgery or pituitary radiation therapy as indicated by growth hormone levels or serum IGF-1 levels that are above the reference range OR
- ii. Patient has severe diarrhea and/or flushing episodes associated with metastatic carcinoid tumors OR
- iii. Patient has profuse watery diarrhea associated with Vasoactive Intestinal Peptide (VIP) secreting tumors OR
- iv. Patient has a diagnosis of dumping syndrome AND the following:
 - a. Patient has tried acarbose or has a documented intolerance, FDA labeled contraindication, or hypersensitivity to acarbose OR
- v. Patient has another indication that is supported in CMS approved compendia for the requested agent AND

2. Patient has responded to and tolerated octreotide for a minimum of 2 weeks AND

3. The dose requested is within the FDA approved or CMS approved compendia dosing for the requested agent and indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Initial approval will be for 6 months, renewal approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria and has responded to therapy AND
2. ONE of the following:
 - A. Patient has a diagnosis of acromegaly AND ONE of the following:
 - i. Decrease in growth hormone level to less than 1 ng/mL OR
 - ii. Normalized serum IGF-1 level (IGF-1 is within reference laboratory range) OR
 - iii. Clinical improvement (e.g. reduction in tumor size, decreased headaches, improved cardiovascular or respiratory symptoms) OR
 - B. Patient has severe diarrhea and/or flushing episodes associated with metastatic carcinoid tumors OR
 - C. Patient has profuse watery diarrhea associated with Vasoactive Intestinal Peptide (VIP) secreting tumors OR
 - D. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
3. The dose requested is within the FDA approved or CMS approved compendia dosing for the requested agent and indication

Prior Authorization Group – Somatostatin Analogs PA - Somatuline Depot

Drug Name(s):

SOMATULINE DEPOT

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Program applies to new starts only.

Criteria for initial approval require ONE of the following:

1. Patient has an FDA approved indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:

A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

B. Prescriber states the patient is currently using the requested agent OR

2. BOTH of the following:

A. ONE of the following

i. Patient has a diagnosis of acromegaly AND ONE of the following:

a. Patient is not a candidate for surgical resection or pituitary radiation therapy OR

b. The requested agent is for adjunctive therapy with pituitary radiation therapy OR

c. Patient had an inadequate response to surgery or pituitary radiation therapy as indicated by growth hormone levels or serum IGF-1 levels that are above the reference range OR

ii. Patient has a diagnosis of gastroenteropancreatic neuroendocrine tumors AND ONE of the following:

a. The tumors are unresectable, locally advanced, well or moderately differentiated OR

b. The tumors have metastasized OR

iii. Patient has a diagnosis of carcinoid syndrome (i.e. flushing and/or diarrhea) OR

iv. Patient has another indication that is supported in CMS approved compendia for the requested agent AND

B. The dose requested is within the FDA approved or CMS approved compendia dosing for the requested agent and indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Initial approval will be for 6 months, renewal approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ONE of the following:

1. Patient has an FDA approved indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently using the requested agent OR
2. Patient has been previously approved for the requested agent through the plan's PA criteria and has responded to therapy AND BOTH of the following:
 - A. ONE of the following:
 - i. Patient has a diagnosis of acromegaly AND the following:
 - a. Patient has had clinical improvement (e.g. reduction in tumor size, decreased headaches, improved cardiovascular or respiratory symptoms) OR
 - ii. Patient has a diagnosis of metastatic OR unresectable, locally advanced, well or moderately differentiated gastroenteropancreatic neuroendocrine tumors OR
 - iii. Patient has a diagnosis of carcinoid syndrome (i.e. flushing and/or diarrhea) OR
 - iv. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
 - B. The dose requested is within the FDA approved or CMS approved compendia dosing for the requested agent and indication

Prior Authorization Group – Somatostatin Analogs PA - Somavert

Drug Name(s):

SOMAVERT

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require BOTH of the following:

1. Patient has a diagnosis of acromegaly AND ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently using the requested agent AND is at risk if therapy is changed OR
 - C. BOTH of the following:
 - A. ONE of the following:
 - a. Patient is not a candidate for surgical resection or pituitary radiation therapy OR
 - b. Patient had an inadequate response to surgery or pituitary radiation therapy as indicated by serum IGF-1 levels that are above the reference range AND
 - B. ONE of the following:
 - a. Patient has tried and failed a prerequisite agent (octreotide or Somatuline Depot) OR
 - b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to prerequisite agents AND
2. The dose requested is within the FDA approved dosing for the requested agent and indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria and has responded to therapy AND
2. Patient has a diagnosis of acromegaly AND the following:
 - A. Patient has had clinical improvement (e.g. reduction in tumor size, decreased headaches, improved cardiovascular or respiratory symptoms) AND
3. The dose requested is within the FDA approved dosing for the requested agent and indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Initial approval will be for 6 months, renewal approval will be for 12 months

Other Criteria:

Prior Authorization Group – Sovaldi PA

Drug Name(s):

SOVALDI

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The dosing of the requested agent is within the FDA labeled dosing or is AASLD/IDSA guideline supported AND
5. ONE of the following:
 - A. There is documentation that the patient is currently being treated with the requested agent OR
 - B. Patient has an FDA labeled contraindication or hypersensitivity to the preferred agents: Harvoni (ledipasvir/sofosbuvir) for supported genotypes OR Epclusa (sofosbuvir/velpatasvir) for genotype 2 or 3 OR
 - C. Prescriber has submitted documentation based on FDA approved labeling or AASLD/IDSA guidelines to support the use of the non-preferred agent for the patient's diagnosis and genotype over the preferred agents: Harvoni (ledipasvir/sofosbuvir) for supported genotypes OR Epclusa (sofosbuvir/velpatasvir) for genotype 2 or 3

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group – Spravato PA

Drug Name(s):

SPRAVATO

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require BOTH of the following:

1. BOTH of following:
 - a. Patient has a diagnosis of treatment-resistant depression (TRD) AND
 - b. Patient is currently receiving treatment with an oral antidepressant and will continue to receive the antidepressant with the requested agent AND
2. ONE of following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent OR
 - c. ALL of the following:
 - i. Patient has had an inadequate response to at least two different oral antidepressants (e.g., SSRIs, SNRIs) AND
 - ii. Prescriber is a specialist (e.g., psychiatrist) or the prescriber has consulted with a specialist AND
 - iii. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. BOTH of the following:
 - a. Patient has a diagnosis of treatment-resistant depression (TRD) AND
 - b. Patient is currently receiving treatment with an oral antidepressant and will continue to receive the antidepressant with the requested agent AND
3. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent OR
 - c. ALL of the following:

- i. Patient has shown clinical benefit with the requested agent AND
- ii. Prescriber is a specialist (e.g., psychiatrist) or the prescriber has consulted with a specialist AND
- iii. Patient does NOT have any FDA labeled contraindication(s) to the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Strensiq PA

Drug Name(s):

STRENSIQ

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has ONE of the following diagnoses:
 - a. Perinatal or infantile-onset hypophosphatasia OR
 - b. Juvenile-onset hypophosphatasia (HPP) AND
2. Patient has clinical manifestations consistent with hypophosphatasia (e.g. vitamin B6-dependent seizures, skeletal abnormalities: such as rachitic chest deformity leading to respiratory problems or bowed arms/legs, “failure to thrive”) AND
3. Patient has radiographic imaging to support the diagnosis of hypophosphatasia (e.g. infantile rickets, alveolar bone loss, craniosynostosis) AND
4. Patient has confirmation of ALPL gene mutation AND
5. Patient has a measured total serum alkaline phosphatase (ALP) level that is below the normal lab reference range for age and sex AND
6. Patient has ONE of the following:
 - a. Elevated urine concentration of phosphoethanolamine (PEA) OR
 - b. Elevated serum concentration of pyridoxal 5'-phosphate (PLP) in the absence of vitamin supplements within one week prior to the test OR
 - c. Elevated urinary inorganic pyrophosphate (PPi)

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has ONE of the following diagnoses:
 - a. Perinatal or infantile-onset hypophosphatasia OR
 - b. Juvenile-onset hypophosphatasia (HPP) AND
3. Patient has responded to treatment with the requested agent as evidenced by an improvement and/or stabilization respiratory status, growth, or radiographic findings

Age Restrictions:

Prescriber Restrictions:

Prescriber is an endocrinologist, a specialist in metabolic or bone disease, or the prescriber has consulted with an endocrinologist or a specialist in metabolic or bone disease

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Substrate Reduction Therapy PA - Miglustat

Drug Name(s):

miglustat 100 mg capsule

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of Gaucher disease type 1 confirmed by ONE of the following:
 - A. A baseline glucocerebrosidase enzyme activity of less than 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR
 - B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND
2. ONE of the following:
 - A. Patient's medication history indicates use of at least ONE enzyme replacement therapy (i.e. Cerezyme, Vpriv, Elelyso) OR
 - B. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to at least ONE enzyme replacement therapy AND
3. Prescriber has drawn baseline levels of hemoglobin, platelets, liver volume, and spleen volume AND
4. Prior to any treatment for the intended diagnosis, the patient has had ONE of the following:
 - A. Anemia (defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender) OR
 - B. Thrombocytopenia (defined as platelet count of less than 100,000 per microliter) OR
 - C. Hepatomegaly OR
 - D. Splenomegaly OR
 - E. Growth failure (i.e. growth velocity is below the standard mean for age) OR
 - F. Evidence of bone disease with other causes ruled out

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of Gaucher disease type 1 AND
3. Patient has shown improvement and/or stabilization from baseline in ONE of the following:

- A. Spleen volume OR
- B. Hemoglobin level OR
- C. Liver volume OR
- D. Platelet count OR
- E. Growth OR
- F. Bone pain or crisis

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. endocrinologist, geneticist, hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Symdeko PA

Drug Name(s):

SYMDEKO

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of cystic fibrosis AND
2. ONE of the following:
 - A. Patient has the presence of the F508del mutation on both alleles (homozygous) of the CFTR gene confirmed by genetic testing OR
 - B. Patient has ONE of the CFTR gene mutations as indicated in the FDA label as confirmed by genetic testing AND
3. Patient has had pre-therapeutic/baseline FEV1 levels measured AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with another CFTR agent [e.g. Kalydeco (ivacaftor), Orkambi (lumacaftor/ivacaftor)] OR
 - B. Patient is currently being treated with another CFTR agent AND will discontinue the other CFTR agent prior to starting the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. Patient has a diagnosis of cystic fibrosis AND
3. Patient has shown improvement or stabilization in FEV1 from pretreatment levels with the requested agent AND
4. ONE of the following:
 - A. Patient is NOT currently being treated with another CFTR agent [e.g. Kalydeco (ivacaftor), Orkambi (lumacaftor/ivacaftor)] OR
 - B. Patient is currently being treated with another CFTR agent AND will discontinue the other CFTR agent prior to starting the requested agent

Age Restrictions:

Patient is 6 years of age or over

Prescriber Restrictions:

Prescriber is a specialist (e.g. cystic fibrosis, pulmonologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Technivie PA

Drug Name(s):

TECHNIVIE

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The dosing of the requested agent is within the FDA labeled dosing or is AASLD/IDSA guideline supported AND
5. ONE of the following:
 - A. There is documentation that the patient is currently being treated with the requested agent OR
 - B. Patient has an FDA labeled contraindication or hypersensitivity to the preferred agent: Harvoni (ledipasvir/sofosbuvir) for supported genotypes OR
 - C. Prescriber has submitted documentation based on FDA approved labeling or AASLD/IDSA guidelines to support the use of the non-preferred agent for the patient's diagnosis and genotype over the preferred agent: Harvoni (ledipasvir/sofosbuvir) for supported genotypes

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group – Tetrabenazine PA

Drug Name(s):

**tetrabenazine tablet
XENAZINE**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. ONE of the following:
 - A. Patient has a diagnosis of chorea associated with Huntington's disease OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
2. If the patient has a current diagnosis of depression, the patient is being treated for depression AND
3. If the patient has a diagnosis of suicidal ideation and/or behavior, the patient must not be actively suicidal AND
4. Patient is NOT receiving a monoamine oxidase inhibitor (MAOI) OR the patient's MAOI will be discontinued at least 14 days before starting therapy with the requested agent AND
5. Patient is NOT receiving reserpine OR the patient's reserpine will be discontinued at least 20 days before starting therapy with the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Topical Doxepin PA

Drug Name(s):

**PRUDOXIN
ZONALON**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require the following:

1. ONE of the following:

a. Patient has a diagnosis of moderate pruritus associated with atopic dermatitis
AND ONE of the following:

i. Patient's medication history includes the use of a topical corticosteroid
(e.g. clobetasol, hydrocortisone, triamcinolone) OR

ii. Patient has a documented intolerance, FDA labeled contraindication, or
hypersensitivity to a topical corticosteroid OR

b. Patient has a diagnosis of moderate pruritus associated with lichen simplex
chronicus AND ONE of the following:

i. Patient's medication history includes the use of a topical corticosteroid
(e.g. clobetasol, hydrocortisone, triamcinolone) OR

ii. Patient has a documented intolerance, FDA labeled contraindication, or
hypersensitivity to a topical corticosteroid

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 8 days

Other Criteria:

Prior Authorization Group – Topical NSAID PA - Voltaren

Drug Name(s):

diclofenac 1% gel

VOLTAREN 1% gel

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. ONE of the following:

a. Patient has an FDA labeled indication for the requested agent OR

b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

a. Patient's medication history includes use of any prescription oral NSAID (non-steroidal anti-inflammatory drug) OR

b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to any prescription oral NSAID

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

3 months for acute pain, 12 months for all other diagnoses

Other Criteria:

Prior Authorization Group – Trelstar PA

Drug Name(s):

**TRELSTAR
TRELSTAR MIXJECT**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

This program applies to new starts only.

Criteria for approval require ALL of the following:

1. ONE of the following:
 - a. Patient has an FDA labeled indication for the requested agent OR
 - b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent OR
 - c. BOTH of the following:
 - i. Patient is NOT currently being treated with the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindication(s) to the requested agent AND
3. The dose requested is within the FDA labeled or CMS approved compendia dosing for the requested indication

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Trientine PA

Drug Name(s):

SYPRINE

trientine capsule

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require BOTH of the following:

1. Patient has a diagnosis of Wilson’s disease confirmed by ONE of the following:
 - a. Confirmation of genetic mutation of the ATP7B gene OR
 - b. Patient has TWO of the following:
 - i. Presence of hepatic abnormality (e.g. acute liver failure, cirrhosis, fatty liver)
 - ii. Presence of Kayser-Fleischer rings
 - iii. Serum ceruloplasmin level less than 20 mg/dL
 - iv. Basal urinary copper excretion greater than 40 mcg/24 hours or the testing laboratory’s upper limit of normal
 - v. Hepatic parenchymal copper content greater than 40 mcg/g dry weight
 - vi. Presence of neurological symptoms (e.g. dystonia, hypertonia, rigidity with tremors, dysarthria, muscle spasms, dysphasia, polyneuropathy, dysautonomia) AND
2. ONE of the following:
 - a. Patient’s medication history indicates use of a penicillamine (e.g. Depen) OR
 - b. Patient has a documented intolerance, FDA labeled contraindication or hypersensitivity to a penicillamine

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan’s PA criteria AND
2. Patient has a diagnosis of Wilson’s disease AND
3. Patient has responded to treatment with the requested agent as evidenced by ONE of the following:
 - a. Improvement and/or stabilization in hepatic abnormality OR
 - b. Reduction in Kayser-Fleischer rings OR

- c. Improvement and/or stabilization in neurological symptoms (e.g. dystonia, hypertonia, rigidity with tremors, dysarthria, muscle spasms, dysphasia, polyneuropathy, dysautonomia) OR
- d. Basal urinary copper excretion greater than 200 mcg/24 hours

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. gastroenterologist, hepatologist, or neurologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Tymlos PA

Drug Name(s):

TYMLOS

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

An increased baseline risk for osteosarcoma

Required Medical:

Criteria for approval require ALL of the following:

1. Patient is a postmenopausal female with a diagnosis of osteoporosis AND ONE of the following:
 - a. Patient has a history of vertebral fracture(s), or low trauma or fragility fracture(s) [e.g. prior fracture from minor trauma such as falling from standing height or less] within the past 5 years OR
 - b. Patient has a T-score that is -2.5 or lower (greater than or equal to 2.5 SD below the mean BMD value for a young adult) AND ONE of the following:
 - i. Patient has failed either a bisphosphonate or SERM OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate or SERM AND
2. ONE of the following:
 - a. Patient is not receiving concomitant bisphosphonate, SERM, Forteo (teriparatide), Prolia (denosumab), or Xgeva (denosumab) therapy within the past 90 days OR
 - b. Prescriber indicates that the patient will discontinue the current bisphosphonate, SERM, Forteo (teriparatide), Prolia (denosumab), or Xgeva (denosumab) therapy prior to initiating therapy with the requested agent AND
3. The dose requested is within the FDA approved labeling AND
4. The total duration of treatment with Tymlos has not exceeded 2 years

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

No prior Tymlos use approve 2 yrs, Prior Tymlos use approve remainder of 2 yrs of total therapy

Other Criteria:

Prior Authorization Group – Urea Cycle Disorders PA - Buphenyl

Drug Name(s):

**BUPHENYL tablet
sodium phenylbutyrate powder, tablet**

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of ONE of the following:
 - a. Urea cycle disorder with neonatal-onset involving deficiencies of carbamylphosphate synthetase, ornithine transcarbamylase, or argininosuccinic acid synthetase OR
 - b. Urea cycle disorder with late-onset and history of hyperammonemic encephalopathy involving deficiencies of carbamylphosphate synthetase, ornithine transcarbamylase, or argininosuccinic acid synthetase AND
2. The dose requested is within the FDA labeled dosing for the requested indication

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. geneticist, metabolic disorders) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Viekira PA

Drug Name(s):

VIEKIRA PAK

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The dosing of the requested agent is within the FDA labeled dosing or is AASLD/IDSA guideline supported AND
5. If genotype 1, the patient's subtype has been identified and provided AND
6. ONE of the following:
 - A. There is documentation that the patient is currently being treated with the requested agent OR
 - B. Patient has an FDA labeled contraindication or hypersensitivity to the preferred agent: Harvoni (ledipasvir/sofosbuvir) for supported genotypes OR
 - C. Prescriber has submitted documentation based on FDA approved labeling or AASLD/IDSA guidelines to support the use of the non-preferred agent for the patient's diagnosis and genotype over the preferred agent: Harvoni (ledipasvir/sofosbuvir) for supported genotypes

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group – Viekira XR PA

Drug Name(s):

VIEKIRA XR

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The dosing of the requested agent is within the FDA labeled dosing or is AASLD/IDSA guideline supported AND
5. If genotype 1, the patient's subtype has been identified and provided AND
6. ONE of the following:
 - A. There is documentation that the patient is currently being treated with the requested agent OR
 - B. Patient has an FDA labeled contraindication or hypersensitivity to the preferred agent: Harvoni (ledipasvir/sofosbuvir) for supported genotypes OR
 - C. Prescriber has submitted documentation based on FDA approved labeling or AASLD/IDSA guidelines to support the use of the non-preferred agent for the patient's diagnosis and genotype over the preferred agent: Harvoni (ledipasvir/sofosbuvir) for supported genotypes

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group – Voriconazole PA

Drug Name(s):

VFEND IV

voriconazole injection, suspension, tablet

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for initial approval require ONE of the following:

- A. Patient has a diagnosis of invasive *Aspergillus* OR
- B. Patient has an infection caused by *Scedosporium apiospermum* or *Fusarium* species OR
- C. Patient has a diagnosis of esophageal candidiasis or candidemia in nonneutropenic patient AND patient has tried fluconazole or an alternative antifungal agent OR patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to fluconazole or an alternative antifungal agent OR
- D. Patient has a diagnosis of blastomycosis AND patient has tried itraconazole OR patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to itraconazole OR
- E. The requested agent is being prescribed for prophylaxis of invasive *Aspergillus* or *Candida* AND patient is severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR
- F. Patient has another indication that is supported in CMS approved compendia for the requested agent

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

One month for esophageal candidiasis, 6 months for other indications

Other Criteria:

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. ONE of the following:
 - A. Patient has a diagnosis of invasive Aspergillus, Scedosporium apiospermum, Fusarium, esophageal candidiasis, candidemia in nonneutropenic patient or blastomycosis and patient has continued indicators of active disease (e.g. continued radiologic findings, positive cultures, positive serum galactomannan assay for Aspergillus) OR
 - B. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or Candida and patient continues to be severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR
 - C. Patient has another indication that is supported in CMS approved compendia for the requested agent

Prior Authorization Group – Vosevi PA

Drug Name(s):

VOSEVI

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The dosing of the requested agent is within the FDA labeled dosing or is AASLD/IDSA guideline supported AND
5. If genotype 1, the patient's subtype has been identified and provided

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group – Vyndamax PA

Drug Name(s):

VYNDAMAX

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND
2. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND
3. Vyndamax (tafamidis) and Vyndaqel (tafamidis meglumine) will not be used concurrently with each other

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND
3. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND
4. Patient has shown clinical benefit with the requested agent AND
5. Vyndamax (tafamidis) and Vyndaqel (tafamidis meglumine) will not be used concurrently with each other

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g., cardiologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Vyndaqel PA

Drug Name(s):

VYNDAQEL

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND
2. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND
3. Vyndamax (tafamidis) and Vyndaqel (tafamidis meglumine) will not be used concurrently with each other

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND
3. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND
4. Patient has shown clinical benefit with the requested agent AND
5. Vyndamax (tafamidis) and Vyndaqel (tafamidis meglumine) will not be used concurrently with each other

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g., cardiologist) or the prescriber has consulted with a specialist

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Xgeva PA

Drug Name(s):

XGEVA

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. ONE of the following:

A. Patient has a diagnosis of multiple myeloma AND ALL of the following:

i. The requested agent will be used for the prevention of skeletal-related events AND

ii. ONE of the following:

1. Patient has tried and failed zoledronic acid (Zometa) OR

2. Patient has a documented intolerance, FDA labeled contraindication or hypersensitivity to zoledronic acid (Zometa) AND

iii. ONE of the following:

1. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR

2. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR

3. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR

B. Patient has a solid tumor cancer diagnosis (e.g. thyroid, non-small cell lung, kidney cancer, prostate cancer, or breast cancer) AND ALL of the following:

i. The requested agent will be used for the prevention of skeletal-related events AND

ii. Patient has bone metastases AND

iii. ONE of the following:

1. Patient has tried and failed zoledronic acid (Zometa) OR

2. Patient has a documented intolerance, FDA labeled contraindication or hypersensitivity to zoledronic acid (Zometa) AND

iv. ONE of the following:

1. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR
2. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR
3. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR

Criteria continues: see Other Criteria

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- C. Patient has a diagnosis of giant cell tumor of bone AND ONE of the following:
 - i. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR
 - ii. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR
 - iii. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR
 - D. Patient has a diagnosis of hypercalcemia of malignancy AND ONE of the following:
 - i. Patient has tried and failed zoledronic acid (Zometa) OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to zoledronic acid (Zometa) AND
2. ONE of the following:
 - A. Patient is NOT receiving concomitant Prolia (denosumab) therapy within the past 90 days OR
 - B. Patient will discontinue the current Prolia (denosumab) therapy prior to initiating therapy with the requested agent AND
 3. The requested dose is within the FDA labeled dosing for the requested indication

Prior Authorization Group – Xolair PA

Drug Name(s):

XOLAIR

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

Required Medical:

Criteria for initial approval require ONE of the following:

1. Patient has a diagnosis of asthma AND ALL of the following:
 - a. If the patient is 6 to less than 12 years of age, the patient meets BOTH of the following:
 - i. Patient's pretreatment IgE level is 30 IU/mL to 1300 IU/mL AND
 - ii. Patient's weight is 20 kg to 150 kg AND
 - b. If the patient is 12 years of age or over, the patient meets ALL of the following:
 - i. Patient's pretreatment IgE level is 30 IU/mL to 700 IU/mL AND
 - ii. Patient's weight is 30 kg to 150 kg AND
 - iii. Patient has a baseline FEV1 less than 80% predicted AND
 - c. Allergic asthma has been confirmed by positive skin test or in vitro reactivity test (RAST) to a perennial aeroallergen AND
 - d. There is evidence of a claim that the patient is currently using an inhaled corticosteroid within the past 90 days OR the patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to inhaled corticosteroid AND
 - e. ONE of the following:
 - i. There is evidence of a claim that the patient is currently using a long-acting beta-2 agonist within the past 90 days OR the patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to long-acting beta-2 agonist OR
 - ii. There is evidence of a claim that the patient is currently using a leukotriene modifier or theophylline within the past 90 days OR the patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to a leukotriene modifier or theophylline AND
 - f. Patient is experiencing exacerbations of asthma symptoms AND
 - g. The requested dose is within the FDA labeled dose OR

Initial criteria continues: see Other Criteria

Age Restrictions:

For diagnosis of asthma, patient is 6 years of age or over. For diagnosis of chronic idiopathic urticaria, patient is 12 years of age or over.

Prescriber Restrictions:

Coverage Duration:

Initial: 24 weeks for asthma and chronic idiopathic urticaria Renewal: 12 months

Other Criteria:

2. Patient has a diagnosis of chronic idiopathic urticaria AND ALL of the following:
 - a. Patient has a history of chronic idiopathic urticaria for at least 6 months AND
 - b. Patient has a history of hives and itching AND
 - c. ONE of the following:
 - i. There is evidence of a claim that the patient is currently on maximum tolerable H1 antihistamine therapy within the past 90 days OR
 - ii. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to H1 antihistamine therapy AND
 - d. The requested dose is within the FDA labeled dose

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's PA criteria AND
2. ONE of the following:
 - a. Patient has a diagnosis of asthma AND ALL of the following:
 - i. Patient's weight is within the FDA indicated range for their age (i.e. 20 kg to 150 kg for patients age 6 to less than 12 years and 30 kg to 150 kg for patients 12 years of age or over) AND
 - ii. Patient does not have clinical worsening defined as ONE of the following:
 1. Increase in inhaled corticosteroid use
 2. Treatment with systemic corticosteroids
 3. Increased use of short acting beta-2 agonist rescue medication
 4. Unscheduled care visits (urgent care, ER, or hospitalizations) due to exacerbations AND
 - iii. ONE of the following:
 1. There is evidence of a claim that the patient is currently on standard therapy (such as a combination of an inhaled corticosteroid, long acting beta-2 agonist, leukotriene receptor antagonist, theophylline, oral corticosteroid or an oral beta-2 agonist tablet) within the past 90 days OR

- 2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ALL standard therapies AND
 - iv. The requested dose is within the FDA labeled dose OR
- b. Patient has a diagnosis of chronic idiopathic urticaria AND BOTH of the following:
 - i. Improvement in symptoms (e.g. number of hives, size of hives, reduction in itching) AND
 - ii. The requested dose is within the FDA labeled dose

Prior Authorization Group – Xyrem PA

Drug Name(s):

XYREM

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ONE of the following:

A. Patient has a diagnosis of narcolepsy with cataplexy OR

B. Patient has a diagnosis of narcolepsy with excessive daytime sleepiness AND BOTH of the following:

i. ONE of the following:

a. Patient is under 18 years of age OR

b. ONE of the following:

1. Patient's medication history indicates the use of modafinil or armodafinil OR

2. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to modafinil or armodafinil AND

ii. ONE of the following:

a. Patient's medication history indicates the use of ONE standard stimulant agent (e.g. methylphenidate) OR

b. Patient has a documented intolerance, FDA labeled contraindication, or hypersensitivity to ONE standard stimulant agent (e.g. methylphenidate)

Age Restrictions:

Patient is 7 years of age or over

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group – Zepatier PA

Drug Name(s):

ZEPATIER

Covered Uses:

All FDA-approved indications not otherwise excluded from Part D. All medically accepted indications not otherwise excluded from Part D.

Exclusion Criteria:

FDA labeled contraindication(s) to the requested agent

Required Medical:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The dosing of the requested agent is within the FDA labeled dosing or is AASLD/IDSA guideline supported AND
5. If genotype 1, the patient's subtype has been identified and provided AND
6. If genotype 1a, the prescriber has tested the patient for the presence of virus with NS5A resistance-associated polymorphisms AND
7. ONE of the following:
 - A. There is documentation that the patient is currently being treated with the requested agent OR
 - B. Patient has an FDA labeled contraindication or hypersensitivity to the preferred agents: Harvoni (ledipasvir/sofosbuvir) for supported genotypes OR Eplclusa (sofosbuvir/velpatasvir) for genotype 3 OR
 - C. Prescriber has submitted documentation based on FDA approved labeling or AASLD/IDSA guidelines to support the use of the non-preferred agent for the patient's diagnosis and genotype over the preferred agents: Harvoni (ledipasvir/sofosbuvir) for supported genotypes OR Eplclusa (sofosbuvir/velpatasvir) for genotype 3

Age Restrictions:

Prescriber Restrictions:

Prescriber is a specialist (e.g. gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria: